

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11) Publication number:

**0 359 472 B1**

(12)

**EUROPEAN PATENT SPECIFICATION**

(45) Date of publication of patent specification: 27.12.95 (51) Int. Cl.<sup>6</sup>: **C12N 15/32, C12P 21/02, A01N 63/00**

(21) Application number: **89309069.6**

(22) Date of filing: **07.09.89**

Divisional application 95201374.6 filed on 07/09/89.

The file contains technical information submitted after the application was filed and not included in this specification

(54) **Synthetic insecticidal crystal protein gene**

(30) Priority: **09.09.88 US 242482**

(43) Date of publication of application: **21.03.90 Bulletin 90/12**

(45) Publication of the grant of the patent: **27.12.95 Bulletin 95/52**

(84) Designated Contracting States:  
**AT BE CH DE ES FR GB GR IT LI LU NL SE**

(56) References cited:  
**EP-A- 0 204 590**

**PROC. NATL. ACAD. SCI. USA, vol. 83, October 1987, pp. 7036-7040, Washington, US; V. SEKAR et al.**

**BIOTECHNOLOGY, vol. 6, January 1988, pp. 61-66, New York, US; S.A. McPHERSON et al.**

**GENET. SEL. EVOL., vol. 19, no. 2, 1987, pp. 143-154, Versailles, FR; M. BOUDRAA**

(73) Proprietor: **Mycogen Plant Science, Inc.**  
**1209 Orange Street**  
**Wilmington**  
**Delaware 19801 (US)**

(72) Inventor: **Adang, Michael J.**  
**160 Tamarack Drive**  
**Athens**  
**Georgia 30605 (US)**  
Inventor: **Rocheleau, Thomas A.**  
**3100 Buena Vista Street**  
**Madison**  
**Wisconsin 53704 (US)**  
Inventor: **Merlo, Donald J.**  
**601 Hampden Court**  
**Midland**  
**Michigan 48640 (US)**  
Inventor: **Murray, Elizabeth E.**  
**2426 Commonwealth Avenue**  
**Madison**  
**Wisconsin 53711 (US)**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

**EP 0 359 472 B1**

NUCLEIC ACIDS RESEARCH, vol. 17, no. 2,  
1989, pp. 477-498, Lada, GB; E.E. MURRAY et  
al.

PROC. NATL. ACAD. SCI. USA, vol. 84, De-  
cember 1987, pp. 8961-8965, Washington, US;  
B.A. SPRINGER et al.

NATURE, vol. 328, 2 July 1987, pp. 33-37,  
London, GB; M. VAECK et al.

PNAS, Vol. 88, 1991, PP. 3324-3328; Perlak et  
al.

⑦ Representative: Fisher, Adrian John et al  
CARPMAELS & RANSFORD  
43 Bloomsbury Square  
London WC1A 2RA (GB)

## Description

## FIELD OF THE INVENTION

5 This invention relates to the field of bacterial molecular biology and, in particular, to genetic engineering by recombinant technology for the purpose of protecting plants from insect pests. Disclosed herein are the chemical synthesis of a modified crystal protein gene from Bacillus thuringiensis var. tenebrionis (Btt), and the selective expression of this synthetic insecticidal gene. Also disclosed is the transfer of the cloned synthetic gene into a host microorganism, rendering the organism capable of producing, at improved levels  
10 of expression, a protein having toxicity to insects. This invention facilitates the genetic engineering of bacteria and plants to attain desired expression levels of novel toxins having agronomic value.

## BACKGROUND OF THE INVENTION

15 B. thuringiensis (Bt) is unique in its ability to produce, during the process of sporulation, proteinaceous, crystalline inclusions which are found to be highly toxic to several insect pests of agricultural importance. The crystal proteins of different Bt strains have a rather narrow host range and hence are used commercially as very selective biological insecticides. Numerous strains of Bt are toxic to lepidopteran and dipteran insects. Recently two subspecies (or varieties) of Bt have been reported to be pathogenic to  
20 coleopteran insects: var. tenebrionis (Krieg et al. (1983) Z. Angew. Entomol. 96:500-508) and var. san diego (Herrnstadt et al. (1986) Biotechnol. 4:305-308). Both strains produce flat, rectangular crystal inclusions and have a major crystal component of 64-68 kDa (Herrnstadt et al. supra; Bernhard (1986) FEMS Microbiol. Lett. 33:261-265).

Toxin genes from several subspecies of Bt have been cloned and the recombinant clones were found to  
25 be toxic to lepidopteran and dipteran insect larvae. The two coleopteran-active toxin genes have also been isolated and expressed. Herrnstadt et al. supra clones a 5.8 kb BamHI fragment of Bt var. san diego DNA. The protein expressed in E. coli was toxic to P. luteola (Elm leaf beetle) and had a molecular weight of approximately 83 kDa. This 83 kDa toxin product from the var. san diego gene was larger than the 64 kDa crystal protein isolated from Bt var. san diego cells, suggesting that the Bt var. san diego crystal protein  
30 may be synthesized as a larger precursor molecule that is processed by Bt var. san diego but not by E. coli prior to being formed into a crystal.

Sekar et al. (1987) Proc. Nat. Acad. Sci. USA 84:7036-7040; U.S. Patent Application 108,285, filed October 13, 1987 isolated the crystal protein gene from Btt and determined the nucleotide sequence. This crystal protein gene was contained on a 5.9 kb BamHI fragment (pNSBF544). A subclone containing the 3  
35 kb HindIII fragment from pNSBF544 was constructed. This HindIII fragment contains an open reading frame (ORF) that encodes a 644-amino acid polypeptide of approximately 73 kDa. Extracts of both subclones exhibited toxicity to larvae of Colorado potato beetle (Leptinotarsa decemlineata, a coleopteran insect). 73- and 65-kDa peptides that cross-reacted with an antiserum against the crystal protein of var. tenebrionis were produced on expression in E. coli. Sporulating var. tenebrionis cells contain an immunoreactive 73-kDa peptide that corresponds to the expected product from the ORF of pNSBP544. However, isolated crystals  
40 primarily contain a 65-kDa component. When the crystal protein gene was shortened at the N-terminal region, the dominant protein product obtained was the 65-kDa peptide. A deletion derivative, p544Pst-Met5, was enzymatically derived from the 5.9 kb BamHI fragment upon removal of forty-six amino acid residues from the N-terminus. Expression of the N-terminal deletion derivative, p544Pst-Met5, resulted in the  
45 production of, almost exclusively, the 65 kDa protein. Recently, McPherson et al. (1988) Biotechnology 6:61-66 demonstrated that the Btt gene contains two functional translational initiation codons in the same reading frame leading to the production of both the full-length protein and an N-terminal truncated form.

Chimeric toxin genes from several strains of Bt have been expressed in plants. Four modified Bt2 genes from var. berliner 1715, under the control of the 2' promoter of the Agrobacterium TR-DNA, were  
50 transferred into tobacco plants (Vaecck et al. (1987) Nature 328:33-37). Insecticidal levels of toxin were produced when truncated genes were expressed in transgenic plants. However, the steady state mRNA levels in the transgenic plants were so low that they could not be reliably detected in Northern blot analysis and hence were quantified using ribonuclease protection experiments. Bt mRNA levels in plants producing the highest level of protein corresponded to  $\approx 0.0001\%$  of the poly(A)<sup>+</sup> mRNA.

55 In the report by Vaecck et al. (1987) supra, expression of chimeric genes containing the entire coding sequence of Bt2 were compared to those containing truncated Bt2 genes. Additionally, some T-DNA constructs included a chimeric NPTII gene as a marker selectable in plants, whereas other constructs carried translational fusions between fragments of Bt2 and the NPTII gene. Insecticidal levels of toxin were

produced when truncated Bt2 genes or fusion constructs were expressed in transgenic plants. Greenhouse grown plants produced  $\approx 0.02\%$  of the total soluble protein as the toxin, or  $3\mu\text{g}$  of toxin per g. fresh leaf tissue and, even at five-fold lower levels, showed 100% mortality in six-day feeding assays. However, no significant insecticidal activity could be obtained using the intact Bt2 coding sequence, despite the fact that the same promoter was used to direct its expression. Intact Bt2 protein and RNA yields in the transgenic plant leaves were 10 - 50 times lower than those for the truncated Bt2 polypeptides or fusion proteins.

Barton *et al.* (1987) *Plant Physiol.* **85**:1103-1109 showed expression of a Bt protein in a system containing a 35S promoter, a viral (TMV) leader sequence, the Bt HD-1 4.5 kb gene (encoding a 645 amino acid protein followed by two proline residues) and a nopaline synthase (*nos*) poly(A)<sup>+</sup> sequence. Under these conditions expression was observed for Bt mRNA at levels up to  $47\text{ pg}/20\mu\text{g}$  RNA and  $12\text{ ng}/\text{mg}$  plant protein. This amount of Bt protein in plant tissue produced 100% mortality in two days. This level of expression still represents a low level of mRNA ( $2.5 \times 10^{-4}\%$ ) and protein ( $1.2 \times 10^{-3}\%$ ).

Various hybrid proteins consisting of N-terminal fragments of increasing length of the Bt2 protein fused to NPTII were produced in *E. coli* by Hofte *et al.* (1988) *FEBS Lett.* **226**:364-370. Fusion proteins containing the first 607 amino acids of Bt2 exhibited insect toxicity; fusion proteins not containing this minimum N-terminal fragment were nontoxic. Appearance of NPTII activity was not dependent upon the presence of insecticidal activity; however, the conformation of the Bt2 polypeptide appeared to exert an important influence on the enzymatic activity of the fused NPTII protein. This study did suggest that the global 3-D structure of the Bt2 polypeptide is disturbed in truncated polypeptides.

A number of researchers have attempted to express plant genes in yeast (Neill *et al.* (1987) *Gene* **55**:303-317; Rothstein *et al.* (1987) *Gene* **55**:353-356; Coraggio *et al.* (1986) *EMBO J.* **5**:459-465) and *E. coli* (Fuzakawa *et al.* (1987) *FEBS Lett.* **224**:125-127; Vies *et al.* (1986) *EMBO J.* **5**:2439-2444; Gatenby *et al.* (1987) *Eur. J. Biochem.* **168**:227-231). In the case of wheat  $\alpha$ -gliadin (Neill *et al.* (1987) *supra*),  $\alpha$ -amylase (Rothstein *et al.* (1987) *supra*) genes, and maize zein genes (Coraggio *et al.* (1986) *supra*) in yeast, low levels of expression have been reported. Neill *et al.* have suggested that the low levels of expression of  $\alpha$ -gliadin in yeast may be due in part to codon usage bias, since  $\alpha$ -gliadin codons for Phe, Leu, Ser, Gly, Tyr and especially Glu do not correlate well with the abundant yeast isoacceptor tRNAs. In *E. coli* however, soybean glycinin A2 (Fuzakawa *et al.* (1987) *supra*) and wheat RuBPC SSU (Vies *et al.* (1986) *supra*; Gatenby *et al.* (1987) *supra*) are expressed adequately.

Not much is known about the makeup of tRNA populations in plants. Viotti *et al.* (1978) *Biochim. Biophys. Acta* **517**:125-132 report that maize endosperm actively synthesizing zein, a storage protein rich in glutamine, leucine, and alanine, is characterized by higher levels of accepting activity for these three amino acids than are maize embryo tRNAs. This may indicate that the tRNA population of specific plant tissues may be adapted for optimum translation of highly expressed proteins such as zein. To our knowledge, no one has experimentally altered codon bias in highly expressed plant genes to determine possible effects of the protein translation in plants to check the effects on the level of expression.

#### SUMMARY OF THE INVENTION

It is the overall object of the present invention to provide a means for plant protection against insect damage. The invention disclosed herein comprises a modified Bt insecticidal protein gene. This gene is designed to be expressed in plants at a level higher than a native Bt gene. It is preferred that the synthetic gene be designed to be highly expressed in plants as defined herein. Preferably, the synthetic gene is at least approximately 85% homologous to an insecticidal protein gene of Bt.

It is a particular object of this invention to provide a synthetic structural gene coding for an insecticidal protein from Btt having, for example, the nucleotide sequences presented in Figure 1 and spanning nucleotides 1 through 1793 or spanning nucleotide 1 through 1833 with functional equivalence.

In designing synthetic Btt genes of this invention for enhanced expression in plants, the DNA sequence of the native Btt structural gene is modified in order to attain an A+T content in nucleotide base composition substantially that found in plants, and also preferably to form a plant initiation sequence, and to eliminate sequences that cause destabilization, inappropriate polyadenylation, degradation and termination of RNA and to avoid sequences that constitute secondary structure hairpins and RNA splice sites. In the synthetic genes, codons used to specify a given amino acid are selected with regard to the distribution frequency of codon usage employed in highly expressed plant genes to specify that amino acid. As is appreciated by those skilled in the art, the distribution frequency of codon usage utilized in the synthetic gene is a determinant of the level of expression. Hence, the synthetic gene is designed such that its distribution frequency of codon usage deviates, preferably, no more than 25% from that of highly expressed plant genes and, more preferably, no more than about 10%. In addition, consideration is given to the

percentage G+C content of the degenerate third base (monocotyledons appear to favor G+C in this position, whereas dicotyledons do not). It is also recognized that the XCG nucleotide is the least preferred codon in dicots whereas the XTA codon is avoided in both monocots and dicots. The synthetic genes of this invention also preferably have CG and TA doublet avoidance indices as defined in the Detailed Description closely approximating those of the chosen host plant. More preferably these indices deviate from that of the host by no more than about 10-15%.

Assembly of the Bt gene of this invention is performed using standard technology known to the art. The Btt structural gene designed for enhanced expression in plants of the specific embodiment is enzymatically assembled within a DNA vector from chemically synthesized oligonucleotide duplex segments. The synthetic Bt gene is then introduced into a plant host cell and expressed by means known to the art. The insecticidal protein produced upon expression of the synthetic Bt gene in plants is functionally equivalent to a native Bt crystal protein in having toxicity to the same insects.

#### BRIEF DESCRIPTION OF THE FIGURES

Figure 1 presents the nucleotide sequence for the synthetic Btt gene. Where different, the native sequence as found in p544Pst-Met5 is shown above. Changes in amino acids (underlined) occur in the synthetic sequence with alanine replacing threonine at residue 2 and leucine replacing the stop at residue 596 followed by the addition of 13-amino acids at the C-terminus.

Figure 2 represents a simplified scheme used in the construction of the synthetic Btt gene. Segments A through M represent oligonucleotide pieces annealed and ligated together to form DNA duplexes having unique splice sites to allow specific enzymatic assembly of the DNA segments to give the desired gene.

Figure 3 is a schematic diagram showing the assembly of oligonucleotide segments in the construction of a synthetic Btt gene. Each segment (A through M) is built from oligonucleotides of different sizes, annealed and ligated to form the desired DNA segment.

#### DETAILED DESCRIPTION OF THE INVENTION

The following definitions are provided in order to provide clarity as to the intent or scope of their usage in the Specification and claims.

Expression refers to the transcription and translation of a structural gene to yield the encoded protein. The synthetic Bt genes of the present invention are designed to be expressed at a higher level in plants than the corresponding native Bt genes. As will be appreciated by those skilled in the art, structural gene expression levels are affected by the regulatory DNA sequences (promoter, polyadenylation sites, enhancers, etc.) employed and by the host cell in which the structural gene is expressed. Comparisons of synthetic Bt gene expression and native Bt gene expression must be made employing analogous regulatory sequences and in the same host cell. It will also be apparent that analogous means of assessing gene expression must be employed in such comparisons.

Promoter refers to the nucleotide sequences at the 5' end of a structural gene which direct the initiation of transcription. Promoter sequences are necessary, but not always sufficient, to drive the expression of a downstream gene. In prokaryotes, the promoter drives transcription by providing binding sites to RNA polymerases and other initiation and activation factors. Usually promoters drive transcription preferentially in the downstream direction, although promotional activity can be demonstrated (at a reduced level of expression) when the gene is placed upstream of the promoter. The level of transcription is regulated by promoter sequences. Thus, in the construction of heterologous promoter/structural gene combinations, the structural gene is placed under the regulatory control of a promoter such that the expression of the gene is controlled by promoter sequences. The promoter is positioned preferentially upstream to the structural gene and at a distance from the transcription start site that approximates the distance between the promoter and the gene it controls in its natural setting. As is known in the art, some variation in this distance can be tolerated without loss of promoter function.

A gene refers to the entire DNA portion involved in the synthesis of a protein. A gene embodies the structural or coding portion which begins at the 5' end from the translational start codon (usually ATG) and extends to the stop (TAG, TGA or TAA) codon at the 3' end. It also contains a promoter region, usually located 5' or upstream to the structural gene, which initiates and regulates the expression of a structural gene. Also included in a gene are the 3' end and poly(A)+ addition sequences.

Structural gene is that portion of a gene comprising a DNA segment encoding a protein, polypeptide or a portion thereof, and excluding the 5' sequence which drives the initiation of transcription. The structural gene may be one which is normally found in the cell or one which is not normally found in the cellular

location wherein it is introduced, in which case it is termed a heterologous gene. A heterologous gene may be derived in whole or in part from any source known to the art, including a bacterial genome or episome, eukaryotic, nuclear or plasmid DNA, cDNA, viral DNA or chemically synthesized DNA. A structural gene may contain one or more modifications in either the coding or the untranslated regions which could affect the biological activity or the chemical structure of the expression product, the rate of expression or the manner of expression control. Such modifications include, but are not limited to, mutations, insertions, deletions and substitutions of one or more nucleotides. The structural gene may constitute an uninterrupted coding sequence or it may include one or more introns, bounded by the appropriate splice junctions. The structural gene may be a composite of segments derived from a plurality of sources, naturally occurring or synthetic. The structural gene may also encode a fusion protein.

Synthetic gene refers to a DNA sequence of a structural gene that is chemically synthesized in its entirety or for the greater part of the coding region. As exemplified herein, oligonucleotide building blocks are synthesized using procedures known to those skilled in the art and are ligated and annealed to form gene segments which are then enzymatically assembled to construct the entire gene. As is recognized by those skilled in the art, functionally and structurally equivalent genes to the synthetic genes described herein may be prepared by site-specific mutagenesis or other related methods used in the art.

Transforming refers to stably introducing a DNA segment carrying a functional gene into an organism that did not previously contain that gene.

Plant tissue includes differentiated and undifferentiated tissues of plants, including but not limited to, roots, shoots, leaves, pollen, seeds, tumor tissue and various forms of cells in culture, such as single cells, protoplasts, embryos and callus tissue. The plant tissue may be in planta or in organ, tissue or cell culture.

Plant cell as used herein includes plant cells in planta and plant cells and protoplasts in culture.

Homology refers to identity or near identity of nucleotide or amino acid sequences. As is understood in the art, nucleotide mismatches can occur at the third or wobble base in the codon without causing amino acid substitutions in the final polypeptide sequence. Also, minor nucleotide modifications (e.g., substitutions, insertions or deletions) in certain regions of the gene sequence can be tolerated and considered insignificant whenever such modifications result in changes in amino acid sequence that do not alter functionality of the final product. It has been shown that chemically synthesized copies of whole, or parts of, gene sequences can replace the corresponding regions in the natural gene without loss of gene function. Homologs of specific DNA sequences may be identified by those skilled in the art using the test of cross-hybridization of nucleic acids under conditions of stringency as is well understood in the art (as described in Hames and Higgins (eds.) (1985) Nucleic Acid Hybridization, IRL Press, Oxford, UK). Extent of homology is often measured in terms of percentage of identity between the sequences compared.

Functionally equivalent refers to identity or near identity of function. A synthetic gene product which is toxic to at least one of the same insect species as a natural Bt protein is considered functionally equivalent thereto. As exemplified herein, both natural and synthetic Bt genes encode 65 kDa, insecticidal proteins having essentially identical amino acid sequences and having toxicity to coleopteran insects. The synthetic Bt genes of the present invention are not considered to be functionally equivalent to native Bt genes, since they are expressible at a higher level in plants than native Bt genes.

Frequency of preferred codon usage refers to the preference exhibited by a specific host cell in usage of nucleotide codons to specify a given amino acid. To determine the frequency of usage of a particular codon in a gene, the number of occurrences of that codon in the gene is divided by the total number of occurrences of all codons specifying the same amino acid in the gene. Table 1, for example, gives the frequency of codon usage for Bt genes, which was obtained by analysis of four Bt genes whose sequences are publicly available. Similarly, the frequency of preferred codon usage exhibited by a host cell can be calculated by averaging frequency of preferred codon usage in a large number of genes expressed by the host cell. It is preferable that this analysis be limited to genes that are highly expressed by the host cell. Table 1, for example, gives the frequency of codon usage by highly expressed genes exhibited by dicotyledonous plants, and monocotyledonous plants. The dicot codon usage was calculated using 154 highly expressed coding sequences obtained from Genbank which are listed in Table 1. Monocot codon usage was calculated using 53 monocot nuclear gene coding sequences obtained from Genbank and listed in Table 1, located in Example 1.

When synthesizing a gene for improved expression in a host cell it is desirable to design the gene such that its frequency of codon usage approaches the frequency of preferred codon usage of the host cell.

The percent deviation of the frequency of preferred codon usage for a synthetic gene from that employed by a host cell is calculated first by determining the percent deviation of the frequency of usage of a single codon from that of the host cell followed by obtaining the average deviation over all codons. As defined herein this calculation includes unique codons (i.e., ATG and TGG). The frequency of preferred

codon usage of the synthetic Bt gene, whose sequence is given in Figure 1, is given in Table 1. The frequency of preferred usage of the codon 'GTA' for valine in the synthetic gene (0.10) deviates from that preferred by dicots (0.12) by  $0.02/0.12 = 0.167$  or 16.7%. The average deviation over all amino acid codons of the Bt synthetic gene codon usage from that of dicot plants is 7.8%. In general terms the overall average deviation of the codon usage of a synthetic gene from that of a host cell is calculated using the equation

$$n=1-Z \frac{\frac{X_n - Y_n}{X_n} \times 100}{Z}$$

where  $X_n$  = frequency of usage for codon n in the host cell;  $Y_n$  = frequency of usage for codon n in the synthetic gene. Where n represents an individual codon that specifies an amino acid, the total number of codons is Z, which in the preferred embodiment is 61. The overall deviation of the frequency of codon usage for all amino acids should preferably be less than about 25%, and more preferably less than about 10%.

Derived from is used to mean taken, obtained, received, traced, replicated or descended from a source (chemical and/or biological). A derivative may be produced by chemical or biological manipulation (including but not limited to substitution, addition, insertion, deletion, extraction, isolation, mutation and replication) of the original source.

Chemically synthesized, as related to a sequence of DNA, means that the component nucleotides were assembled in vitro. Manual chemical synthesis of DNA may be accomplished using well established procedures (Caruthers, M. (1983) in Methodology of DNA and RNA Sequencing, Weissman (ed.), Praeger Publishers, New York, Chapter 1); or automated chemical synthesis can be performed using one of a number of commercially available machines.

The term, designed to be highly expressed as used herein refers to a level of expression of a designed gene wherein the amount of its specific mRNA transcripts produced is sufficient to be quantified in Northern blots and, thus, represents a level of specific mRNA expressed corresponding to greater than or equal to approximately 0.001% of the poly(A)+ mRNA. To date, natural Bt genes are transcribed at a level wherein the amount of specific mRNA produced is insufficient to be estimated using the Northern blot technique. However, in the present invention, transcription of a synthetic Bt gene designed to be highly expressed not only allows quantification of the specific mRNA transcripts produced but also results in enhanced expression of the translation product which is measured in insecticidal bioassays.

Crystal protein or insecticidal crystal protein or crystal toxin refers to the major protein component of the parasporal crystals formed in strains of Bt. This protein component exhibits selective pathogenicity to different species of insects. The molecular size of the major protein isolated from parasporal crystals varies depending on the strain of Bt from which it is derived. Crystal proteins having molecular weights of approximately 132, 65, and 28 kDa have been reported. It has been shown that the approximately 132 kDa protein is a protoxin that is cleaved to form an approximately 65 kDa toxin.

The crystal protein gene refers to the DNA sequence encoding the insecticidal crystal protein in either full length protoxin or toxin form, depending on the strain of Bt from which the gene is derived.

The authors of this invention observed that expression in plants of Bt crystal protein mRNA occurs at levels that are not routinely detectable in Northern blots and that low levels of Bt crystal protein expression correspond to this low level of mRNA expression. It is preferred for exploitation of these genes as potential biocontrol methods that the level of expression of Bt genes in plant cells be improved and that the stability of Bt mRNA in plants be optimized. This will allow greater levels of Bt mRNA to accumulate and will result in an increase in the amount of insecticidal protein in plant tissues. This is essential for the control of insects that are relatively resistant to Bt protein.

Thus, this invention is based on the recognition that expression levels of desired, recombinant insecticidal protein in transgenic plants can be improved via increased expression of stabilized mRNA transcripts; and that, conversely, detection of these stabilized RNA transcripts may be utilized to measure expression of translational product (protein). This invention provides a means of resolving the problem of low expression of insecticidal protein RNA in plants and, therefore, of low protein expression through the use of an improved, synthetic gene specifying an insecticidal crystal protein from Bt.



Attempts to improve the levels of expression of Bt genes in plants have centered on comparative studies evaluating parameters such as gene type, gene length, choice of promoters, addition of plant viral untranslated RNA leader, addition of intron sequence and modification of nucleotides surrounding the initiation ATG codon. To date, changes in these parameters have not led to significant enhancement of Bt protein expression in plants. Applicants find that, surprisingly, to express Bt proteins at the desired level in plants, modifications in the coding region of the gene were effective. Structural-function relationships can be studied using site-specific mutagenesis by replacement of restriction fragments with synthetic DNA duplexes containing the desired nucleotide changes (Lo et al. (1984) Proc. Natl. Acad. Sci. 81:2285-2289). However, recent advances in recombinant DNA technology now make it feasible to chemically synthesize an entire gene designed specifically for a desired function. Thus, the Btt coding region was chemically synthesized, modified in such a way as to improve its expression in plants. Also, gene synthesis provides the opportunity to design the gene so as to facilitate its subsequent mutagenesis by incorporating a number of appropriately positioned restriction endonuclease sites into the gene.

The present invention provides a synthetic Bt gene for a crystal protein toxic to an insect. As exemplified herein, this protein is toxic to coleopteran insects. To the end of improving expression of this insecticidal protein in plants, this invention provides a DNA segment homologous to a Btt structural gene and, as exemplified herein, having approximately 85% homology to the Btt structural gene in p544Pst-Met5. In this embodiment the structural gene encoding a Btt insecticidal protein is obtained through chemical synthesis of the coding region. A chemically synthesized gene is used in this embodiment because it best allows for easy and efficacious accommodation of modifications in nucleotide sequences required to achieve improved levels of cross-expression.

Today, in general, chemical synthesis is a preferred method to obtain a desired modified gene. However, to date, no plant protein gene has been chemically synthesized nor has any synthetic gene for a bacterial protein been expressed in plants. In this invention, the approach adopted for synthesizing the gene consists of designing an improved nucleotide sequence for the coding region and assembling the gene from chemically synthesized oligonucleotide segments. In designing the gene, the coding region of the naturally-occurring gene, preferably from the Btt subclone, p544Pst-Met5, encoding a 65 kDa polypeptide having coleoperan toxicity, is scanned for possible modifications which would result in improved expression of the synthetic gene in plants. For example, to optimize the efficiency of translation, codons preferred in highly expressed proteins of the host cell are utilized.

Bias in codon choice within genes in a single species appears related to the level of expression of the protein encoded by that gene. Codon bias is most extreme in highly expressed proteins of E. coli and yeast. In these organisms, a strong positive correlation has been reported between the abundance of an isoaccepting tRNA species and the favored synonymous codon. In one group of highly expressed proteins in yeast, over 96% of the amino acids are encoded by only 25 of the 61 available codons (Bennetzen and Hall (1982) J. Biol. Chem. 257:3026-3031).

These 25 codons are preferred in all sequenced yeast genes, but the degree of preference varies with the level of expression of the genes. Recently, Hoekema and colleagues (1987) Mol. Cell. Biol. 7:2914-2924 reported that replacement of these 25 preferred codons by minor codons in the 5' end of the highly expressed yeast gene PGK1 results in a decreased level of both protein and mRNA. They concluded that biased codon choice in highly expressed genes enhances translation and is required for maintaining mRNA stability in yeast. Without doubt, the degree of codon bias is an important factor to consider when engineering high expression of heterologous genes in yeast and other systems.

Experimental evidence obtained from point mutations and deletion analysis has indicated that in eukaryotic genes specific sequences are associated with post-transcriptional processing, RNA destabilization, translational termination, intron splicing and the like. These are preferably employed in the synthetic genes of this invention. In designing a bacterial gene for expression in plants, sequences which interfere with the efficacy of gene expression are eliminated.

In designing a synthetic gene, modifications in nucleotide sequence of the coding region are made to modify the A + T content in DNA base composition of the synthetic gene to reflect that normally found in genes for highly expressed proteins native to the host cell. Preferably the A + T content of the synthetic gene is substantially equal to that of said genes for highly expressed proteins. In genes encoding highly expressed plant proteins, the A + T content is approximately 55%. It is preferred that the synthetic gene have an A + T content near this value, and not sufficiently high as to cause destabilization of RNA and, therefore, lower the protein expression levels. Thus, the A + T content is no more than 60% and most preferably is about 55%. Also, for ultimate expression in plants, the synthetic gene nucleotide sequence is preferably modified to form a plant initiation sequence at the 5' end of the coding region. In addition, particular attention is preferably given to assure that unique restriction sites are placed in strategic positions

to allow efficient assembly of oligonucleotide segments during construction of the synthetic gene and to facilitate subsequent nucleotide modification. As a result of these modifications in coding region of the native Bt gene, the preferred synthetic gene is expressed in plants at an enhanced level when compared to that observed with natural Bt structural genes.

5 In specific embodiments, the synthetic Bt gene of this invention encodes a Btt protein toxic to coleopteran insects. Preferably, the toxic polypeptide is about 598 amino acids in length, is at least 75% homologous to a Btt polypeptide, and, as exemplified herein, is essentially identical to the protein encoded by p544Pst-Met5, except for replacement of threonine by alanine at residue 2. This amino acid substitution results as a consequence of the necessity to introduce a guanine base at position +4 in the coding  
10 sequence.

In designing the synthetic gene of this invention, the coding region from the Btt subclone, p544Pst-Met5, encoding a 65 kDa polypeptide having coleopteran toxicity, is scanned for possible modifications which would result in improved expression of the synthetic gene in plants. For example, in preferred  
15 embodiments, the synthetic insecticidal protein is strongly expressed in dicot plants, e.g., tobacco, tomato, cotton, etc., and hence, a synthetic gene under these conditions is designed to incorporate to advantage codons used preferentially by highly expressed dicot proteins. In embodiments where enhanced expression of insecticidal protein is desired in a monocot, codons preferred by highly expressed monocot proteins (given in Table 1) are employed in designing the synthetic gene.

In general, genes within a taxonomic group exhibit similarities in codon choice, regardless of the  
20 function of these genes. Thus an estimate of the overall use of the genetic code by a taxonomic group can be obtained by summing codon frequencies of all its sequenced genes. This species-specific codon choice is reported in this invention from analysis of 208 plant genes. Both monocot and dicot plants are analyzed individually to determine whether these broader taxonomic groups are characterized by different patterns of synonymous codon preference. The 208 plant genes included in the codon analysis code for proteins  
25 having a wide range of functions and they represent 6 monocot and 36 dicot species. These proteins are present in different plant tissues at varying levels of expression.

In this invention it is shown that the relative use of synonymous codons differs between the monocots and the dicots. In general, the most important factor in discriminating between monocot and dicot patterns of codon usage is the percentage G + C content of the degenerate third base. In monocots, 16 of 18 amino  
30 acids favor G + C in this position, while dicots only favor G + C in 7 of 18 amino acids.

The G ending codons for Thr, Pro, Ala and Ser are avoided in both monocots and dicots because they contain C in codon position II. The CG dinucleotide is strongly avoided in plants (Boudraa (1987) Genet. Sel. Evol. 19:143-154) and other eukaryotes (Grantham et al. (1985) Bull. Inst. Pasteur 83:95-148), possibly due to regulation involving methylation. In dicots, XCG is always the least favored codon, while in monocots  
35 this is not the case. The doublet TA is also avoided in codon positions II and III in most eukaryotes, and this is true of both monocots and dicots.

Grantham and colleagues (1986) Oxford Surveys in Evol. Biol. 3:48-81 have developed two codon choice indices to quantify CG and TA doublet avoidance in codon positions II and III. XCG/XCC is the ratio of codons having C as base II of G-ending to C-ending triplets, while XTA/XTT is the ratio of A-ending to T-  
40 ending triplets with T as the second base. These indices have been calculated for the plant data in this paper (Table 2) and support the conclusion that monocot and dicot species differ in their use of these dinucleotides.

45

50

55

Table 2

Avoidance of CG and TA doublets in codons position II-III.  
XCG/XCC and XTA/XAA values are multiplied by 100.

Group	Plants	Dicots	Mono- cots	Maize	Soy- bean	RuBPC SSU	CAB
XCG/XCC	40	30	61	67	37	18	22
XTA/XTT	37	35	47	43	41	9	13

RuBPC SSU = ribulose 1,5 biphosphate small subunit  
CAB = chlorophyll a/b binding protein

Additionally, for two species, soybean and maize, species-specific codon usage profiles were calculated (not shown). The maize codon usage pattern resembles that of monocots in general, since these sequences represent over half of the monocot sequences available. The codon profile of the maize subsample is even more strikingly biased in its preference for G+C in codon position III. On the other hand, the soybean codon usage pattern is almost identical to the general dicot pattern, even though it represents a much smaller portion of the entire dicot sample.

In order to determine whether the coding strategy of highly expressed genes such as the ribulose 1,5 biphosphate small subunit (RuBPC SSU) and chlorophyll a/b binding protein (CAB) is more biased than that of plant genes in general, codon usage profiles for subsets of these genes (19 and 17 sequences, respectively) were calculated (not shown). The RuBPC SSU and CAB pooled samples are characterized by stronger avoidance of the codons XCG and XTA than in the larger monocot and dicot samples (Table 2). Although most of the genes in these subsamples are dicot in origin (17/19 and 15/17), their codon profile resembles that of the monocots in that G+C is utilized in the degenerate base III.

The use of pooled data for highly expressed genes may obscure identification of species-specific patterns in codon choice. Therefore, the codon choices of individual genes for RuBPC SSU and CAB were tabulated. The preferred codons of the maize and wheat genes for RuBPC SSU and CAB are more restricted in general than are those of the dicot species. This is in agreement with Matsuoka *et al.* (1987) J. Biochem. 102:673-676) who noted the extreme codon bias of the maize RuBPC SSU gene as well as two other highly expressed genes in maize leaves, CAB and phosphoenolpyruvate carboxylase. These genes almost completely avoid the use of A+T in codon position III, although this codon bias was not as pronounced in non-leaf proteins such as alcohol dehydrogenase, zein 22 kDa sub-unit, sucrose synthetase and ATP/ADP translocator. Since the wheat SSU and CAB genes have a similar pattern of codon preference, this may reflect a common monocot pattern for these highly expressed genes in leaves. The CAB gene for Lemna and the RuBPC SSU genes for Chlamydomonas share a similar extreme preference for G+C in codon position III. In dicot CAB genes, however, A+T degenerate bases are preferred by some synonymous codons (e.g., GCT for Ala, CTT for Leu, GGA and GGT for Gly). In general, the G+C preference is less pronounced for both RuBPC SSU and CAB genes in dicots than in monocots.

In designing a synthetic gene for expression in plants, attempts are also made to eliminate sequences which interfere with the efficacy of gene expression. Sequences such as the plant polyadenylation signals, e.g., AATAAA, polymerase II termination sequence, e.g., CAN<sub>7-9</sub>AGTNNAA, UCUUCGG hairpins and plant consensus splice sites are highlighted and, if present in the native *Btt* coding sequence, are modified so as to eliminate potentially deleterious sequences.

Modifications in nucleotide sequence of the *Btt* coding region are also preferably made to reduce the A+T content in DNA base composition. The *Btt* coding region has an A+T content of 64%, which is about 10% higher than that found in a typical plant coding region. Since A+T-rich regions typify plant intergenic regions and plant regulatory regions, it is deemed prudent to reduce the A+T content. The synthetic *Btt* gene is preferably designed to have an A+T content of 55%, in keeping with values usually found in plants.

Also, a single modification (to introduce guanine in lieu of adenine) at the fourth nucleotide position in the Btt coding sequence is made in the preferred embodiment to form a sequence consonant with that believed to function as a plant initiation sequence (Taylor *et al.* (1987) *Mol. Gen. Genet.* 210:572-577) in optimization of expression. In addition, in exemplifying this invention thirty-nine nucleotides (thirteen codons) are added to the coding region of the synthetic gene in an attempt to stabilize primary transcripts. However, it appears that equally stable transcripts are obtained in the absence of this extension polypeptide containing thirty-nine nucleotides.

Not all of the above-mentioned modifications of the natural Bt gene must be made in constructing a synthetic Bt gene in order to obtain enhanced expression. For example, a synthetic gene may be synthesized for other purposes in addition to that of achieving enhanced levels of expression. Under these conditions, the original sequence of the natural Bt gene may be preserved within a region of DNA corresponding to one or more, but not all, segments used to construct the synthetic gene. Depending on the desired purpose of the gene, modification may encompass substitution of one or more, but not all, of the oligonucleotide segments used to construct the synthetic gene by a corresponding region of natural Bt sequence.

As is known to those skilled in the art of synthesizing genes (Mandecki *et al.* (1985) *Proc. Natl. Acad. Sci.* 82:3543-3547; Feretti *et al.* (1986) *Proc. Natl. Acad. Sci.* 83:599-603), the DNA sequence to be synthesized is divided into segment lengths which can be synthesized conveniently and without undue complication. As exemplified herein, in preparing to synthesize the Btt gene, the coding region is divided into thirteen segments (A - M). Each segment has unique restriction sequences at the cohesive ends. Segment A, for example, is 228 base pairs in length and is constructed from six oligonucleotide sections, each containing approximately 75 bases. Single-stranded oligonucleotides are annealed and ligated to form DNA segments. The length of the protruding cohesive ends in complementary oligonucleotide segments is four to five residues. In the strategy evolved for gene synthesis, the sites designed for the joining of oligonucleotide pieces and DNA segments are different from the restriction sites created in the gene.

In the specific embodiment, each DNA segment is cloned into a pC-20 vector for amplification of the DNA. The nucleotide sequence of each fragment is determined at this stage by the dideoxy method using the recombinant phage DNA as templates and selected synthetic oligonucleotides as primers.

As exemplified herein and illustrated schematically in Figures 3 and 4, each segment individually (e.g., segment M) is excised at the flanking restriction sites from its cloning vector and spliced into the vector containing segment A. Most often, segments are added as a paired segment instead of as a single segment to increase efficiency. Thus, the entire gene is constructed in the original plasmid harboring segment A. The nucleotide sequence of the entire gene is determined and found to correspond exactly to that shown in Figure 1.

In preferred embodiments the synthetic Btt gene is expressed in plants at an enhanced level when compared to that observed with natural Btt structural genes. To that end, the synthetic structural gene is combined with a promoter functional in plants, the structural gene and the promoter region being in such position and orientation with respect to each other that the structural gene can be expressed in a cell in which the promoter region is active, thereby forming a functional gene. The promoter regions include, but are not limited to, bacterial and plant promoter regions. To express the promoter region/structural gene combination, the DNA segment carrying the combination is contained by a cell. Combinations which include plant promoter regions are contained by plant cells, which, in turn, may be contained by plants or seeds. Combinations which include bacterial promoter regions are contained by bacteria, e.g., Bt or E. coli. Those in the art will recognize that expression in types of micro-organisms other than bacteria may in some circumstances be desirable and, given the present disclosure, feasible without undue experimentation.

The recombinant DNA molecule carrying a synthetic structural gene under promoter control can be introduced into plant tissue by any means known to those skilled in the art. The technique used for a given plant species or specific type of plant tissue depends on the known successful techniques. As novel means are developed for the stable insertion of foreign genes into plant cells and for manipulating the modified cells, skilled artisans will be able to select from known means to achieve a desired result. Means for introducing recombinant DNA into plant tissue include, but are not limited to, direct DNA uptake (Paszowski, J. *et al.* (1984) *EMBO J.* 3:2717), electroporation (Fromm, M. *et al.* (1985) *Proc. Natl. Acad. Sci. USA* 82:5824), microinjection (Crossway, A. *et al.* (1986) *Mol. Gen. Genet.* 202:179), or T-DNA mediated transfer from Agrobacterium tumefaciens to the plant tissue. There appears to be no fundamental limitation of T-DNA transformation to the natural host range of Agrobacterium. Successful T-DNA-mediated transformation of monocots (Hooykaas-Van Slogteren, G. *et al.* (1984) *Nature* 311:763), gymnosperm (Dandekar, A. *et al.* (1987) *Biotechnology* 5:587) and algae (Ausich, R., EPO application 108,580) has been reported. Representative T-DNA vector systems are described in the following references: An, G. *et al.* (1985) *EMBO*

J. 4:277; Herrera-Estrella, L. et al. (1983) Nature 303:209; Herrera-Estrella, L. et al. (1983) EMBO J. 2:987; Herrera-Estrella, L. et al. (1985) in Plant Genetic Engineering, New York: Cambridge University Press, p. 63. Once introduced into the plant tissue, the expression of the structural gene may be assayed by any means known to the art, and expression may be measured as mRNA transcribed or as protein synthesized.

5 Techniques are known for the *in vitro* culture of plant tissue, and in a number of cases, for regeneration into whole plants. Procedures for transferring the introduced expression complex to commercially useful cultivars are known to those skilled in the art.

In one of its preferred embodiments the invention disclosed herein comprises expression in plant cells of a synthetic insecticidal structural gene under control of a plant expressible promoter, that is to say, by  
10 inserting the insecticide structural gene into T-DNA under control of a plant expressible promoter and introducing the T-DNA containing the insert into a plant cell using known means. Once plant cells expressing a synthetic insecticidal structural gene under control of a plant expressible promoter are obtained, plant tissues and whole plants can be regenerated therefrom using methods and techniques well-known in the art. The regenerated plants are then reproduced by conventional means and the introduced  
15 genes can be transferred to other strains and cultivars by conventional plant breeding techniques.

The introduction and expression of the synthetic structural gene for an insecticidal protein can be used to protect a crop from infestation with common insect pests. Other uses of the invention, exploiting the properties of other insecticide structural genes introduced into other plant species will be readily apparent to those skilled in the art. The invention in principle applies to introduction of any synthetic insecticide  
20 structural gene into any plant species into which foreign DNA (in the preferred embodiment T-DNA) can be introduced and in which said DNA can remain stably replicated. In general, these taxa presently include, but are not limited to, gymnosperms and dicotyledonous plants, such as sunflower (family Compositae), tobacco (family Solanaceae), alfalfa, soybeans and other legumes (family Leguminosae), cotton (family Malvaceae), and most vegetables, as well as monocotyledonous plants. A plant containing in its tissues  
25 increased levels of insecticidal protein will control less susceptible types of insect, thus providing advantage over present insecticidal uses of *Bt*. By incorporation of the insecticidal protein into the tissues of a plant, the present invention additionally provides advantage over present uses of insecticides by eliminating instances of nonuniform application and the costs of buying and applying insecticidal preparations to a field. Also, the present invention eliminates the need for careful timing of application of such preparations since  
30 small larvae are most sensitive to insecticidal protein and the protein is always present, minimizing crop damage that would otherwise result from preapplication larval foraging.

This invention combines the specific teachings of the present disclosure with a variety of techniques and expedients known in the art. The choice of expedients depends on variables such as the choice of insecticidal protein from a *Bt* strain, the extent of modification in preferred codon usage, manipulation of  
35 sequences considered to be destabilizing to RNA or sequences prematurely terminating transcription, insertions of restriction sites within the design of the synthetic gene to allow future nucleotide modifications, addition of introns or enhancer sequences to the 5' and/or 3' ends of the synthetic structural gene, the promoter region, the host in which a promoter region/structural gene combination is expressed, and the like. As novel insecticidal proteins and toxic polypeptides are discovered, and as sequences responsible for  
40 enhanced cross-expression (expression of a foreign structural gene in a given host) are elucidated, those of ordinary skill will be able to select among those elements to produce "improved" synthetic genes for desired proteins having agronomic value. The fundamental aspect of the present invention is the ability to synthesize a novel gene coding for an insecticidal protein, designed so that the protein will be expressed at an enhanced level in plants, yet so that it will retain its inherent property of insect toxicity and retain or  
45 increase its specific insecticidal activity.

## EXAMPLES

The following Examples are presented as illustrations of embodiments of the present invention. They do  
50 not limit the scope of this invention, which is determined by the claims.

The following strains were deposited with the Patent Culture Collection, Northern Regional Research Center, 1815 N. University Street, Peoria, Illinois 61604.

55

Strain	Deposited on	Accession #
<i>E. coli</i> MC1061 (p544-HindIII)	6 October 1987	NRRL B-18257
<i>E. coli</i> MC1061 (p544Pst-Met5)	6 October 1987	NRRL B-18258

The deposited strains are provided for the convenience of those in the art, and are not necessary to practice the present invention, which may be practiced with the present disclosure in combination with publicly available protocols, information, and materials. *E. coli* MC1061, a good host for plasmid transformations, was disclosed by Casadaban, M.J. and Cohen, S.N. (1980) *J. Mol. Biol.* **138**:179-207.

#### Example 1: Design of the synthetic insecticidal crystal protein gene.

##### (i) Preparation of toxic subclones of the Bt gene

Construction, isolation, and characterization of pNSBP544 is disclosed by Sekar, V. et al. (1987) *Proc. Natl. Acad. Sci. USA* **84**:7036-7040, and Sekar, V. and Adang, M.J., U.S. patent application serial no. 108,285, filed October 13, 1987, which is hereby incorporated by reference. A 3.0 kbp *Hind*III fragment carrying the crystal protein gene of pNSBP544 is inserted into the *Hind*III site of pIC-2OH (Marsh, J.L. et al. (1984) *Gene* **32**:481-485), thereby yielding a plasmid designated p544-*Hind*III, which is on deposit. Expression in *E. coli* yields a 73 kDa crystal protein in addition to the 65 kDa species characteristic of the crystal protein obtained from Bt isolates.

A 5.9 kbp *Bam*HI fragment carrying the crystal protein gene is removed from pNSBP544 and inserted into *Bam*HI-linearized pIC-2OH DNA. The resulting plasmid, p405/44-7, is digested with *Bgl*II and religated, thereby removing *Bacillus* sequences flanking the 3'-end of the crystal protein gene. The resulting plasmid, p405/54-12, is digested with *Pst*I and religated, thereby removing *Bacillus* sequences flanking the 5'-end of the crystal protein and about 150 bp from the 5'-end of the crystal protein structural gene. The resulting plasmid, p405/81-4, is digested with *Sph*I and *Pst*I and is mixed with and ligated to a synthetic linker having the following structure:

SD                      MetThrAla  
5' CAGGATCCAACAATGACTGCA3'  
3' GTACGTCCTAGGTTGTTACTG5'  
*Sph*I                      *Pst*I

(SD indicates the location of a Shine-Dalgarno prokaryotic ribosome binding site.) The resulting plasmid, p544Pst-Met5, contains a structural gene encoding a protein identical to one encoded by pNSBP544 except for a deletion of the amino-terminal 47 amino acid residues. The nucleotide sequence of the Bt coding region in p544Pst-Met5 is presented in Figure 1. In bioassays (Sekar and Adang, U.S. patent application serial no. 108,285, *supra*), the proteins encoded by the full-length Bt gene in pNSBP544 and the N-terminal deletion derivative, p544Pst-Met5, were shown to be equally toxic. All of the plasmids mentioned above have their crystal protein genes in the same orientation as the *lacZ* gene of the vector.

##### (ii) Modification of preferred codon usage

Table 1 presents the frequency of codon usage for (A) dicot proteins, (B) Bt proteins, (C) the synthetic Bt gene, and (D) monocot proteins. Although some codons for a particular amino acid are utilized to approximately the same extent by both dicot and Bt proteins (e.g., the codons for serine), for the most part, the distribution of codon frequency varies significantly between dicot and Bt proteins, as illustrated in columns A and B in Table 1.

Table 1. Frequency of Codon Usage

5	Amino Acid	Codon	<u>Distribution Fraction</u>			
			<u>(A)Dicot Genes</u>	<u>(B)Bt Genes</u>	<u>(C)Synthetic Btt Gene</u>	<u>(D)Monocot Genes</u>
10	Gly	GGG	0.12	0.08	0.13	0.21
	Gly	GGA	0.37	0.53	0.37	0.18
	Gly	GGT	0.35	0.24	0.34	0.21
	Gly	GGC	0.16	0.16	0.16	0.40
15	Glu	GAG	0.52	0.13	0.52	0.77
	Glu	GAA	0.48	0.87	0.48	0.23
	Asp	GAT	0.57	0.68	0.56	0.31
	Asp	GAC	0.43	0.32	0.44	0.69
20	Val	GTG	0.30	0.15	0.30	0.38
	Val	GTA	0.12	0.32	0.10	0.07
	Val	GTT	0.38	0.29	0.35	0.20
	Val	GTC	0.20	0.24	0.25	0.34
25	Ala	GCG	0.05	0.12	0.06	0.20
	Ala	GCA	0.26	0.50	0.24	0.16
	Ala	GCT	0.42	0.32	0.41	0.28
	Ala	GCC	0.28	0.06	0.29	0.36
30	Lys	AAG	0.61	0.13	0.58	0.87
	Lys	AAA	0.39	0.87	0.42	0.13
	Asn	AAT	0.45	0.79	0.44	0.23
	Asn	AAC	0.55	0.21	0.56	0.77
35	Met	ATG	1.00	1.00	1.00	1.00
	Ile	ATA	0.19	0.30	0.20	0.09
	Ile	ATT	0.44	0.57	0.43	0.27
	Ile	ATC	0.36	0.13	0.37	0.64
40	Thr	ACG	0.07	0.14	0.07	0.18
	Thr	ACA	0.27	0.68	0.27	0.14
	Thr	ACT	0.36	0.14	0.34	0.22
	Thr	ACC	0.31	0.05	0.32	0.47
45	Trp	TGG	1.00	1.00	1.00	1.00
	End	TGA	0.46	0.00	0.00	0.34
	Cys	TGT	0.43	0.33	0.33	0.27
	Cys	TGC	0.57	0.67	0.67	0.73
50	End	TAG	0.18	0.00	0.00	0.44
	End	TAA	0.37	1.00	1.00	0.22
	Tyr	TAT	0.42	0.81	0.43	0.19
	Tyr	TAC	0.58	0.19	0.57	0.81

55

Table 1 (CONTINUED)

			Distribution Fraction			
5	Amino Acid	Codon	(A) Dicot Genes	(B) Bt Genes	(C) Synthetic Btt. Gene	(D) Monocot Genes
	Phe	TTT	0.45	0.75	0.44	0.28
10	Phe	TTC	0.55	0.25	0.56	0.72
	Ser	AGT	0.14	0.25	0.13	0.07
	Ser	AGC	0.18	0.13	0.19	0.25
	Ser	TCG	0.05	0.08	0.06	0.13
	Ser	TCA	0.18	0.19	0.17	0.13
15	Ser	TCT	0.26	0.25	0.27	0.18
	Ser	TCC	0.19	0.10	0.17	0.24
	Arg	AGG	0.22	0.09	0.23	0.28
	Arg	AGA	0.31	0.50	0.32	0.08
20	Arg	CGG	0.04	0.14	0.05	0.14
	Arg	CGA	0.09	0.14	0.09	0.04
	Arg	CGT	0.23	0.09	0.23	0.11
	Arg	CGC	0.11	0.05	0.09	0.36
25	Gln	CAG	0.38	0.18	0.39	0.43
	Gln	CAA	0.62	0.82	0.61	0.57
	His	CAT	0.52	0.90	0.50	0.38
	His	CAC	0.48	0.10	0.50	0.62
30	Leu	TTG	0.26	0.08	0.27	0.15
	Leu	TTA	0.10	0.46	0.12	0.04
	Leu	CTG	0.09	0.04	0.10	0.27
	Leu	CTA	0.08	0.21	0.10	0.11
35	Leu	CTT	0.29	0.15	0.18	0.16
	Leu	CTC	0.19	0.06	0.22	0.27
	Pro	CCG	0.07	0.20	0.08	0.20
	Pro	CCA	0.44	0.56	0.44	0.39
40	Pro	CCT	0.32	0.24	0.32	0.19
	Pro	CCC	0.16	0.00	0.16	0.22

Bt coding sequences publicly available and 88 coding sequences of dicot nuclear genes were used to compile the codon usage table. The pooled dicot coding sequences, obtained from Genbank, were:



Table 1 (CONTINUED)

5

	GENUS/ SPECIES	GENBANK	PROTEIN	REF
	<i>Antirrhinum majus</i>	AMACHS	Chalcone synthase	
10	<i>Arabidopsis thaliana</i>	ATH1ADH	Alcohol dehydrogenase	
		ATH1H3GA	Histone 3 gene 1	
		ATH1H3GB	Histone 3 gene 2	
		ATH1H4GA	Histone 4 gene 1	
		ATH1H1CP1	CAB	
		ATH1TUBA	$\alpha$ tubulin	
15			5-enolpyruvylshikate 3-phosphate synthase	1
	<i>Benthamia exzelsa</i>		High methionine storage protein	2
	<i>Brassica campestris</i>		Acyl carrier protein	3
	<i>Brassica napus</i>	BNANAP	Napin	
	<i>Brassica oleracea</i>	BOLSLSGR	S-locus specific glycoprotein	
	<i>Canavalia ensiformis</i>	CENCONA	Concanavalin A	
	<i>Carica papaya</i>	CPAPAP	Papain	
20	<i>Chlamydomonas reinhardtii</i>	CRECSS2	Protophyochrome	
		CRERBCS1	RuBPC small subunit gene 1	
		CRERBCS2	RuBPC small subunit gene 2	
	<i>Cucurbita pepo</i>	CUCPIT	Phytochrome	
	<i>Cucumis sativus</i>	CUSGMS	Glyoxosomal malate synthase	
25		CUSLHCPA	CAB	
		CUSSSU	RuBPC small subunit	
	<i>Daucus carota</i>	DAREXT	Extensin	
		DAREXTR	33 kD extensin related protein	
	<i>Dolichos biflorus</i>	DBILECS	seed lectin	
	<i>Flaveria trinervia</i>	FTRBCR	RuBPC small subunit	
30	<i>Glycine max</i>	SOY7SAA	7S storage protein	
		SOYACT1G	Actin 1	
		SOYCTIM	Cl protease inhibitor	
		SOYGLYA1A	Glycinin A1a Bx subunits	
		SOYGLYAA.B	Glycinin ASA4B3 subunits	
		SOYGLYAB	Glycinin A3/b4 subunits	
		SOYGLYR	Glycinin A2B1a subunits	
35		SOYHSP175	Low M W heat shock proteins	
		SOYLGBI	Leghemoglobin	
		SOYLEA	Lectin	
		SOYLOX	Lipoxygenase 1	
		SOYNOD20G	20 kDa nodulin	
		SOYNOD23G	23 kDa nodulin	
		SOYNOD24II	24 kDa nodulin	
40		SOYNOD26B	26 kDa nodulin	
		SOYNOD26R	26 kDa nodulin	
		SOYNOD27R	27 kDa nodulin	
		SOYNOD35M	35 kDa nodulin	
		SOYNOD75	75 kDa nodulin	
		SOYNODR1	Nodulin C51	
45		SOYNODR2	Nodulin E27	
		SOYTRP1	Proline rich protein	
		SOYRUBP	RuBPC small subunit	
		SOYURA	Urease	
		SOYHSP26A	Heat shock protein 26A	4
			Nuclear-encoded chloroplast heat shock protein	5
50			22 kDa nodulin	6
			$\beta$ 1 tubulin	6
			$\beta$ 2 tubulin	6

55

Table 1 (CONTINUED)

	GENUS/ SPECIES	GENBANK	PROTEIN	REF
5	<i>Gossypium hirsutum</i>		Seed $\alpha$ globulin (vicilin)	7
			Seed $\beta$ globulin (vicilin)	7
	<i>Helianthus annuus</i>	HNNRUBCS	RuBPC small subunit	8
	<i>Ipomoea batatas</i>		2S albumin seed storage protein	9
	<i>Lemna gibba</i>	LG1AB19	Wound-induced catalase	
		LG1RS111C	CAB	
10	<i>Lupinus luteus</i>	LUPLBR	RuBPC small subunit	
	<i>Lycopersicon esculentum</i>		leghemoglobin I	
		TOMBI0BR	Biotin binding protein	
		TOMETIYDR	Ethylene biosynthesis protein	
		TOMPG2AR	Polygalacturonase-2a	
		TOMPSI	Tomato photosystem I protein	
		TOMRBCSA	RuBPC small subunit	
15		TOMRBCSB	RuBPC small subunit	
		TOMRBCSC	RuBPC small subunit	
		TOMRBCSD	RuBPC small subunit	
		TOMRRD	Ripening related protein	
		TOMWIPIC	Wound induced proteinase inhibitor I	
		TOMWIPII	Wound induced proteinase inhibitor II	
20			CAB 1A	10
			CAB 1B	10
			CAB 3C	10
			CAB 4	11
			CAB 5	11
25	<i>Medicago sativa</i>	ALFLNJR	Leghemoglobin III	
	<i>Mesembryanthemum crystallinum</i>		RuBPC small subunit	12
	<i>Nicotiana glauca</i>			
	<i>Nicotiana glauca</i>	TOBATT21	Mitochondrial ATP synthase $\beta$ subunit	
			Nitrate reductase	13
			Glutamine synthetase	14
30	<i>Nicotiana tabacum</i>	TOBECI	Endochitinase	
		TOBGAPA	A subunit of chloroplast G3PD	
		TOBGAPB	B subunit of chloroplast G3PD	
		TOBGAPC	C subunit of chloroplast G3PD	
		TOBPR1AR	Pathogenesis related protein 1a	
		TOBPR1CR	Pathogenesis-related protein 1c	
35		TOBPR1PR	Pathogenesis related protein 1b	
		TOBPR1DLF	Peroxidase	
		TOBRBPCO	RuBPC small subunit	
		TOBTIAUR	TMV-induced protein homologous to thaumatin	
	<i>Persea americana</i>	AVOCEL	Cellulase	
40	<i>Peucedanum officinale</i>	PIIOCHL	Chalcone synthase	
	<i>Petunia sp.</i>	PETCAB13	CAB 13	
		PETCAB22L	CAB 22L	
		PETCAB22R	CAB 22R	
		PETCAB25	CAB 25	
		PETCAB37	CAB 37	
45		PETCAB91R	CAB 91R	
		PETCHSR	Chalcone synthase	
		PETGCR1	Glycine-rich protein	
		PETRBCS08	RuBPC small subunit	
		PETRBCS11	RuBPC small subunit	
			70 kDa heat shock protein	15
50	<i>Phaseolus vulgaris</i>	PIIVCHM	Chitinase	
		PIIVDLECA	Phytohemagglutinin E	
		PIIVDLECB	Phytohemagglutinin L	
		PIIVGSR1	Glutamine synthetase 1	
		PIIVGSR2	Glutamine synthetase 2	
55				

Table 1 (CONTINUED)

5	GENUS/ SPECIES	GENBANK	PROTEIN	REF
		PIHVLJA	Leghemoglobin	
		PIHVLCT	Lectin	
		PIHVPAL	Phenylalanine ammonia lyase	
		PIHVTIASAR	$\alpha$ phaseolin	
		PIHVTIASUR	$\beta$ phaseolin	
10	<i>Pisum sativum</i>	PEAALB2	Arcelin seed protein	16
		PEACAB80	Chalcone synthase	17
		PEAGSR1	Seed albumin	
		PEALECA	CAB	
		PEALEGA	Glutamine synthetase (nodule)	
15		PEARUBPS	Lectin	
		PEAVIC2	Legumin	
		PEAVIC4	RuBPC small subunit	
		PEAVIC7	Vicilin	
			Vicilin	
			Alcohol dehydrogenase 1	18
			Glutamine synthetase (leaf)	19
20			Glutamine synthetase (root)	19
			Histone 1	20
			Nuclear encoded chloroplast	4
			heat shock protein	
	<i>Raphanus sativus</i>		RuBPC small subunit	21
	<i>Ricinus communis</i>	RCCAGG	Agglutinin	
25		RCCRICIN	Ricin	
		RCCICL4	Isocitrate lyase	
	<i>Silene pratensis</i>	SIPFDX	Ferrodoxin precursor	
		SIPPCY	Plastocyanin precursor	
	<i>Sinapis alba</i>	SALGAPDII	Nuclear gene for G3PD	
	<i>Solanum tuberosum</i>	POTPAT	Pataatin	
30		POTINHVI	Wound-induced proteinase inhibitor	
		POTLSIG	Light-inducible tissue specific	
			ST-LS1 gene	
		POTPI2G	Wound-induced proteinase inhibitor II	
		POTRBCS	RuBPC small subunit	
35			Sucrose synthase	22
	<i>Spinacia oleracea</i>	SPIACP1	Acyl carrier protein I	
		SPIOEC16	16 kDa photosynthetic oxygen-evolving protein	
		SPIOEC23	23 kDa photosynthetic oxygen-evolving protein	
40		SPIPCG	Plastocyanin	
		SPIPS33	33 kDa photosynthetic water oxidation complex precursor	
			Glycolate oxidase	23
	<i>Vicia faba</i>	VFALBA	Leghemoglobin	
		VFALEB4	Legumin B	
45			Vicillin	24

50 Pooled 53 monocot coding sequences obtained from Genbank (release 55) or, when no Genbank file name is specified, directly from the published source, were:

55

Table 1 (CONTINUED)

5

10

15

20

25

30

35

40

45

50

55

GENUS/SPECIES	GENDANK	PROTEIN	REF
<i>Avena sativa</i>	ASTAP3R	Phytochrome 3	
<i>Hordeum vulgare</i>	BLYALR	Aleurin	
	BLYAMY1	$\alpha$ amylase 1	
	BLYAMY2	$\alpha$ amylase 2	
	BLYCHORD1	Hordein C	
	BLYGLUCB	$\beta$ glucanase	
	BLYHORB	B1 hordein	
	BLYTAPI	Amylase/protease inhibitor	
	BLYTHIAR	Toxin $\alpha$ hordothionin	
	BLYUBIQR	Ubiquitin	
		Histone 3	25
		Leaf specific thionin 1	26
		Leaf specific thionin 2	26
		Plastocyanin	27
<i>Oryza sativa</i>	RICGLUTG	Glutelin	
		Glutelin	28
<i>Triticum aestivum</i>	WHTAMYA	$\alpha$ amylase	
	WHTCAB	CAB	
	WHTEMR	Em protein	
	WHTGIR	gibberellin responsive protein	
	WHTGLGB	$\gamma$ gliadin	
	WHTGLIABA	$\alpha/\beta$ gliadin Class AII	
	WHTGLUTI	High MW glutenin	
	WHTH3	Histone 3	
	WHTH4091	Histone 4	
	WHTRBCB	RuBPC small subunit	
<i>Secale cereale</i>	RYESECGR	$\gamma$ secalin	
<i>Zea mays</i>	MZEAI1G	40.1 kD A1 protein (NADPH-dependent reductase)	
	MZEACTIG	Actin	
	MZEADH11F	Alcohol dehydrogenase 1	
	MZEADH2NR	Alcohol dehydrogenase 2	
	MZEALD	Aldolase	
	MZEANT	ATP/ADP translocator	
	MZEAG2R	Glutelin 2	
	MZEAGST3B	Glutathione S transferase	
	MZEHIJ2	Histone 3	
	MZEHI4C14	Histone 4	
	MZEHS1701	70 kD Heat shock protein, exon 1	
	MZEHS1702	70 kD Heat shock protein, exon 2	
	MZEIJICP	CAB	
	MZEMPL3	Lipid body surface protein L3	
	MZEPEPCR	Phosphoenolpyruvate carboxylase	
	MZERBCS	RuBPC small subunit	
	MZESUSYSG	Sucrose synthetase	
	MZETP12	Triosephosphate isomerase 1	
	MZEZE20M	19 kD zein	
	MZEZE20N	19 kD zein	
	MZEZE15A3	15 kD zein	
	MZEZE16	16 kD zein	
	MZEZE19A	19 kD zein	
	MZEZE22A	22 kD zein	
	MZEZE22B	22 kD zein	
		Catalase 2	29
		Regulatory C1 locus	30

Table 1 (CONTINUED)

Bt codons were obtained from analysis of coding sequences of the following genes: Bt var. kurstaki HD-73, 6.6kb HindIII fragment (Kronstad et al. (1983) J. Bacteriol. 154:419-428); Bt var. kurstaki HD-1, 5.3 kb fragment (Adang et al. (1987) in Biotechnology in Invertebrate Pathology and Cell Culture, K. Maramorosh (ed.), Academic Press, Inc. New York, pp. 85-99); Bt var. kurstaki HD-1, 4.5 kb fragment (Schnepf and Whiteley (1985) J. Biol. Chem. 260:6273-6280); and Bt var. tenebrionis, 3.0 kb HindIII fragment (Sekar et al. (1987) Proc. Natl. Acad. Sci. 84:7036-7040).

# REFERENCES

1. Klee, H.J. et al. (1987) Mol. Gen. Genet. 210:437-442.
2. Altenbach, S.B. et al. (1987) Plant Mol. Biol. 8:239-250.
3. Rose, R.E. et al. (1987) Nucl. Acids Res. 15:7197.
4. Vierling, E. et al. (1988) EMBO J. 7:575-581.
5. Sandal, N.N. et al. (1987) Nucl. Acids Res. 15:1507-1519.
6. Tingey, S.V. et al. (1987) EMBO J. 6:1-9.
7. Chlan, C.A. et al. (1987) Plant Mol. Biol. 9:533-546.
8. Allen, R.D. et al. (1987) Mol. Gen. Genet. 210:211-218.
9. Sakajo, S. et al. (1987) Eur. J. Biochem. 165:437-442.
10. Pirersky, E. et al. (1987) Plant Mol. Biol. 9:109-120.
11. Ray, J. et al. (1987) Nucl. Acids Res. 15:10587.
12. DeRocjer, E.J. et al. (1987) Nucl. Acids Res. 15:6301.
13. Calza, R. et al. (1987) Mol. Gen. Genet. 209:552-562.
14. Tingey, S.V. and Coruzzi, G.M. (1987) Plant Phys. 84:366-373.
15. Winter, J. et al. (1988) Mol. Gen. Genet. 211:315-319.
16. Osborn, T.C. et al. (1988) Science 240:207-210.

Table 1 (CONTINUED)

- 5 17. Ryder, T.B. et al. (1987) Mol. Gen. Genet. 210:219-233.
18. Llewellyn, D.J. et al. (1987) J. Mol. Biol. 195:115-123.
- 10 19. Tingey, S.V. et al. (1987) EMBO J. 6:1-9.
20. Gantt, J.S. and Key, J.L. (1987) Eur. J. Biochem. 166:119-125.
- 15 21. Guidet, F. and Fourcroy, P. (1988) Nucl. Acids Res. 16:2336.
22. Salanoubat, M. and Belliard, G. (1987) Gene 60:47-56.
- 20 23. Volokita, M. and Somerville, C.R. (1987) J. Biol. Chem. 262:15825-15828.
24. Bassner, R. et al. (1987) Nucl. Acids Res. 15:9609.
- 25 25. Chojecki, J. (1986) Carlsberg Res. Commun. 51:211-217.
26. Bohlmann, H. and Apel, K. (1987) Mol. Gen. Genet. 207:446-454.
- 30 27. Nielsen, P.S. and Gausing, K. (1987) FEBS Lett. 225:159-162.
28. Higuchi, W. and Fukazawa, C. (1987) Gene 55:245-253.
- 35 29. Bethards, L.A. et al. (1987) Proc. Natl. Acad. Sci. USA 84:6830-6834.
30. Paz-Ares, J. et al. (1987) EMBO J. 6:3553-3558.

40 For example, dicots utilize the AAG codon for lysine with a frequency of 61% and the AAA codon with a frequency of 39%. In contrast, in Bt proteins the lysine codons AAG and AAA are used with a frequency of 13% and 87%, respectively. It is known in the art that seldom used codons are generally detrimental to that system and must be avoided or used judiciously. Thus, in designing a synthetic gene encoding the Btt crystal protein, individual amino acid codons found in the original Btt gene are altered to reflect the codons preferred by dicot genes for a particular amino acid. However, attention is given to maintaining the overall distribution of codons for each amino acid within the coding region of the gene. For example, in the case of alanine, it can be seen from Table 1 that the codon GCA is used in Bt proteins with a frequency of 50%, whereas the codon GCT is the preferred codon in dicot proteins. In designing the synthetic Btt gene, not all codons for alanine in the original Bt gene are replaced by GCT; instead, only some alanine codons are changed to GCT while others are replaced with different alanine codons in an attempt to preserve the overall distribution of codons for alanine used in dicot proteins. Column C in Table 1 documents that this goal is achieved; the frequency of codon usage in dicot proteins (column A) corresponds very closely to that used in the synthetic Btt gene (column C).

55 In similar manner, a synthetic gene coding for insecticidal crystal protein can be optimized for enhanced expression in monocot plants. In Table 1, column D, is presented the frequency of codon usage of highly expressed monocot proteins.

Because of the degenerate nature of the genetic code, only part of the variation contained in a gene is expressed in this protein. It is clear that variation between degenerate base frequencies is not a neutral

phenomenon since systematic codon preferences have been reported for bacterial, yeast and mammalian genes. Analysis of a large group of plant gene sequences indicates that synonymous codons are used differently by monocots and dicots. These patterns are also distinct from those reported for *E. coli* yeast and man.

5 In general, the plant codon usage pattern more closely resembles that of man and other higher eukaryotes than unicellular organisms, due to the overall preference for G + C content in codon position III. Monocots in this sample share the most commonly used codon for 13 of 18 amino acids as that reported for a sample of human genes (Grantham *et al.* (1986 *supra*), although dicots favor the most commonly used human codon in only 7 of 18 amino acids.

10 Discussions of plant codon usage have focused on the differences between codon choice in plant nuclear genes and in chloroplasts. Chloroplasts differ from higher plants in that they encode only 30 tRNA species. Since chloroplasts have restricted their tRNA genes, the use of preferred codons by chloroplast-encoded proteins appears more extreme. However, a positive correlation has been reported between the level of isoaccepting tRNA for a given amino acid and the frequency with which this codon is used in the chloroplast genome (Pfitzinger *et al.* (1987) *Nucl. Acids Res.* 15:1377-1386).

15 Our analysis of the plant genes sample confirms earlier reports that the nuclear and chloroplast genomes in plants have distinct coding strategies. The codon usage of monocots in this sample is distinct from chloroplast usage, sharing the most commonly used codon for only 1 of 18 amino acids. Dicots in this sample share the most commonly used codon of chloroplasts in only 4 of 18 amino acids. In general, the chloroplast codon profile more closely resembles that of unicellular organisms, with a strong bias towards the use of A + T in the degenerate third base.

In unicellular organisms, highly expressed genes use a smaller subset of codons than do weakly expressed genes although the codons preferred are distinct in some cases. Sharp and Li (1986) *Nucl. Acids Res.* 14:7734-7749 report that codon usage in 165 *E. coli* genes reveals a positive correlation between high expression and increased codon bias. Bennetzen and Hall (1982) *supra* have described a similar trend in codon selection in yeast. Codon usage in these highly expressed genes correlates with the abundance of isoaccepting tRNAs in both yeast and *E. coli*. It has been proposed that the good fit of abundant yeast and *E. coli* mRNA codon usage to isoacceptor tRNA abundance promotes high translation levels and high steady state levels of these proteins. This strongly suggests that the potential for high levels of expression of plant genes in yeast or *E. coli* is limited by their codon usage. Hoekema *et al.* (1987) *supra* report that replacement of the 25 most favored yeast codons with rare codons in the 5' end of the highly expressed gene PGK1 leads to a decrease in both mRNA and protein. These results indicate that codon bias should be emphasized when engineering high expression of foreign genes in yeast and other systems.

### 35 (iii) Sequences within the Btt coding region having potentially destabilizing influences

Analysis of the Btt gene reveals that the A + T content represents 64% of the DNA base composition of the coding region. This level of A + T is about 10% higher than that found in a typical plant coding region. Most often, high A + T regions are found in intergenic regions. Also, many plant regulatory sequences are observed to be AT-rich. These observations lead to the consideration that an elevated A + T content within the Btt coding region may be contributing to a low expression level in plants. Consequently, in designing a synthetic Btt gene, the A + T content is decreased to more closely approximate the A + T levels found in plant proteins. As illustrated in Table 3, the A + T content is lowered to a level in keeping with that found in coding regions of plant nuclear genes. The synthetic Btt gene of this invention has an A + T content of 55%.

Table 3

Adenine + Thymine Content in Btt Coding Region						
Coding Region	Base				%G + C	%A + T
	G	A	T	C		
Natural Btt gene	341	633	514	306	36	64
Synthetic Btt gene	392	530	483	428	45	55

In addition, the natural Btt gene is scanned for sequences that are potentially destabilizing to Btt RNA. These sequences, when identified in the original Btt gene, are eliminated through modification of nucleotide

sequences. Included in this group of potentially destabilizing sequences are:

- (a) plant polyadenylation signals (as described by Joshi (1987) Nucl. Acids Res. 15:9627-9640). In eukaryotes, the primary transcripts of nuclear genes are extensively processed (steps including 5' - capping, intron splicing, polyadenylation) to form mature and translatable mRNAs. In higher plants, polyadenylation involves endonucleolytic cleavage at the polyA site followed by the addition of several A residues to the cleaved end. The selection of the polyA site is presumed to be cis-regulated. During expression of Bt protein and RNA in different plants, the present inventors have observed that the polyadenylated mRNA isolated from these expression systems is not full-length but instead is truncated or degraded. Hence, in the present invention it was decided to minimize possible destabilization of RNA through elimination of potential polyadenylation signals within the coding region of the synthetic Bt gene. Plant polyadenylation signals including AATAAA, AATGAA, AATAAT, AATATT, GATAAA, GATAAA, and AATAAG motifs do not appear in the synthetic Bt gene when scanned for 0 mismatches of the sequences.
- (b) polymerase II termination sequence, CAN<sub>7-9</sub>AGTNAA. This sequence was shown (Vankan and Filipowicz (1988) EMBO J. 7:791-799) to be next to the 3' end of the coding region of the U2 snRNA genes of *Arabidopsis thaliana* and is believed to be important for transcription termination upon 3' end processing. The synthetic Bt gene is devoid of this termination sequence.
- (c) CUUCGG hairpins, responsible for extraordinarily stable RNA secondary structures associated with various biochemical processes (Tuerk et al. (1988) Proc. Natl. Acad. Sci. 85:1364-1368). The exceptional stability of CUUCGG hairpins suggests that they have an unusual structure and may function in organizing the proper folding of complex RNA structures. CUUCGG hairpin sequences are not found with either 0 or 1 mismatches in the Bt coding region.
- (d) plant consensus splice sites, 5' = AAG:GTAAGT and 3' = TTTT(Pu)TTT(Pu)T(Pu)T(Pu)T(Pu)-TGCAG:C, as described by Brown et al. (1986) EMBO J. 5:2749-2758. Consensus sequences for the 5' and 3' splice junctions have been derived from 20 and 30 plant intron sequences, respectively. Although it is not likely that such potential splice sequences are present in Bt genes, a search was initiated for sequences resembling plant consensus splice sites in the synthetic Bt gene. For the 5' splice site, the closest match was with three mismatches. This gave 12 sequences of which two had G:GT. Only position 948 was changed because 1323 has the KpnI site needed for reconstruction. The 3'-splice site is not found in the synthetic Bt gene.

Thus, by highlighting potential RNA-destabilizing sequences, the synthetic Bt gene is designed to eliminate known eukaryotic regulatory sequences that effect RNA synthesis and processing.

## Example 2. Chemical synthesis of a modified Bt structural gene

### (i) Synthesis Strategy

The general plan for synthesizing linear double-stranded DNA sequences coding for the crystal protein from Bt is schematically simplified in Figure 2. The optimized DNA coding sequence (Figure 1) is divided into thirteen segments (segments A-M) to be synthesized individually, isolated and purified. As shown in Figure 2, the general strategy begins by enzymatically joining segments A and M to form segments AM to which is added segment BL to form segment ABLM. Segment CK is then added enzymatically to make segment ABCKLM which is enlarged through addition of segments DJ, EI and RFH sequentially to give finally the total segment ABCDEFGHIJKLM, representing the entire coding region of the Bt gene.

Figure 3 outlines in more detail the strategy used in combining individual DNA segments in order to effect the synthesis of a gene having unique restriction sites integrated into a defined nucleotide sequence. Each of the thirteen segments (A to M) has unique restriction sites at both ends, allowing the segment to be strategically spliced into a growing DNA polymer. Also, unique sites are placed at each end of the gene to enable easy transfer from one vector to another.

The thirteen segments (A to M) used to construct the synthetic gene vary in size. Oligonucleotide pairs of approximately 75 nucleotides each are used to construct larger segments having approximately 225 nucleotide pairs. Figure 3 documents the number of base pairs contained within each segment and specifies the unique restriction sites bordering each segment. Also, the overall strategy to incorporate specific segments at appropriate splice sites is detailed in Figure 3.



(ii) Preparation of oligodeoxynucleotides

Preparation of oligodeoxynucleotides for use in the synthesis of a DNA sequence comprising a gene for Btt is carried out according to the general procedures described by Matteucci et al. (1981) J. Am. Chem. Soc. 103:3185-3192 and Beaucage et al. (1981) Tetrahedron Lett. 22:1859-1862. All oligonucleotides are prepared by the solid-phase phosphoramidite triester coupling approach, using an Applied Biosystems Model 380A DNA synthesizer. Deprotection and cleavage of the oligomers from the solid support are carried out according to standard procedures. Crude oligonucleotide mixtures are purified using an oligonucleotide purification cartridge (OTC, Applied Biosystems) as described by McBride et al. (1988) Biotechniques 6:362-367.

5'-phosphorylation of oligonucleotides is performed with T4 polynucleotide kinase. The reaction contains 2 µg oligonucleotide and 18.2 units polynucleotide kinase (Pharmacia) in linker kinase buffer (Maniatis (1982) Cloning Manual, Fritsch and Sambrook (eds.), Cold Spring Harbor Laboratory, Cold Spring Harbor, NY). The reaction is incubated at 37 °C for 1 hour.

Oligonucleotides are annealed by first heating to 95 °C for 5 min. and then allowing complementary pairs to cool slowly to room temperature. Annealed pairs are reheated to 65 °C, solutions are combined, cooled slowly to room temperature and kept on ice until used. The ligated mixture may be purified by electrophoresis through a 4% NuSieve agarose (FMC) gel. The band corresponding to the ligated duplex is excised, the DNA is extracted from the agarose and ethanol precipitated.

Ligations are carried out as exemplified by that used in M segment ligations. M segment DNA is brought to 65 °C for 25 min, the desired vector is added and the reaction mixture is incubated at 65 °C for 15 min. The reaction is slow cooled over 1-1/2 hours to room temperature. ATP to 0.5mM and 3.5 units of T4 DNA ligase salts are added and the reaction mixture is incubated for 2 hr at room temperature and then maintained overnight at 15 °C. The next morning, vectors which had not been ligated to M block DNA were removed upon linearization by EcoRI digestion. Vectors ligated to the M segment DNA are used to transform E. coli MC1061. Colonies containing inserted blocks are identified by colony hybridization with <sup>32</sup>P-labelled oligonucleotide probes. The sequence of the DNA segment is confirmed by isolating plasmid DNA and sequencing using the dideoxy method of Sanger et al. (1977) Proc. Natl. Acad. Sci. 74:5463-5467.

(iii) Synthesis of Segment AM

Three oligonucleotide pairs (A1 and its complementary strand A1c, A2 and A2C and A3 and A3C) are assembled and ligated as described above to make up segment A. The nucleotide sequence of segment A is as follows:

5  
10  
15  
20  
25  
30  
35  
40

In Table 4, bold lines demarcate the individual oligonucleotides. Fragment A1 contains 71 bases, A1c has 76 bases, A2 has 75 bases, A2C has 76 bases, A3 has 82 bases and A3C has 76 bases. In all, segment A is composed of 228 base pairs and is contained between EcoRI restriction enzyme site and one destroyed EcoRI site (5')J. (Additional restriction sites within Segment A are indicated.) The EcoRI single-stranded cohesive ends allow segment A to be annealed and then ligated to the EcoRI-cut cloning vector, pIC20K.

Segment M comprises three oligonucleotide pairs: M1, 80 bases, M1c, 86 bases, M2, 87 bases, M2c, 87 bases, M3, 85 bases and M3c 79 bases. The individual oligonucleotides are annealed and ligated according to standard procedures as described above. The overall nucleotide sequence of segment M is:

HindIII BspXII  
 EcoRI BclI  
 RI-end  
 1  
 60  
 121  
 181  
 241  
 301  
 361  
 421  
 481  
 541  
 601  
 661  
 721  
 781  
 841  
 901  
 961  
 1021  
 1081  
 1141  
 1201  
 1261  
 1321  
 1381  
 1441  
 1501  
 1561  
 1621  
 1681  
 1741  
 1801  
 1861  
 1921  
 1981  
 2041  
 2101  
 2161  
 2221  
 2281  
 2341  
 2401  
 2461  
 2521  
 2581  
 2641  
 2701  
 2761  
 2821  
 2881  
 2941  
 3001  
 3061  
 3121  
 3181  
 3241  
 3301  
 3361  
 3421  
 3481  
 3541  
 3601  
 3661  
 3721  
 3781  
 3841  
 3901  
 3961  
 4021  
 4081  
 4141  
 4201  
 4261  
 4321  
 4381  
 4441  
 4501  
 4561  
 4621  
 4681  
 4741  
 4801  
 4861  
 4921  
 4981  
 5041  
 5101  
 5161  
 5221  
 5281  
 5341  
 5401  
 5461  
 5521  
 5581  
 5641  
 5701  
 5761  
 5821  
 5881  
 5941  
 6001  
 6061  
 6121  
 6181  
 6241  
 6301  
 6361  
 6421  
 6481  
 6541  
 6601  
 6661  
 6721  
 6781  
 6841  
 6901  
 6961  
 7021  
 7081  
 7141  
 7201  
 7261  
 7321  
 7381  
 7441  
 7501  
 7561  
 7621  
 7681  
 7741  
 7801  
 7861  
 7921  
 7981  
 8041  
 8101  
 8161  
 8221  
 8281  
 8341  
 8401  
 8461  
 8521  
 8581  
 8641  
 8701  
 8761  
 8821  
 8881  
 8941  
 9001  
 9061  
 9121  
 9181  
 9241  
 9301  
 9361  
 9421  
 9481  
 9541  
 9601  
 9661  
 9721  
 9781  
 9841  
 9901  
 9961  
 10021  
 10081  
 10141  
 10201  
 10261  
 10321  
 10381  
 10441  
 10501  
 10561  
 10621  
 10681  
 10741  
 10801  
 10861  
 10921  
 10981  
 11041  
 11101  
 11161  
 11221  
 11281  
 11341  
 11401  
 11461  
 11521  
 11581  
 11641  
 11701  
 11761  
 11821  
 11881  
 11941  
 12001  
 12061  
 12121  
 12181  
 12241  
 12301  
 12361  
 12421  
 12481  
 12541  
 12601  
 12661  
 12721  
 12781  
 12841  
 12901  
 12961  
 13021  
 13081  
 13141  
 13201  
 13261  
 13321  
 13381  
 13441  
 13501  
 13561  
 13621  
 13681  
 13741  
 13801  
 13861  
 13921  
 13981  
 14041  
 14101  
 14161  
 14221  
 14281  
 14341  
 14401  
 14461  
 14521  
 14581  
 14641  
 14701  
 14761  
 14821  
 14881  
 14941  
 15001  
 15061  
 15121  
 15181  
 15241  
 15301  
 15361  
 15421  
 15481  
 15541  
 15601  
 15661  
 15721  
 15781  
 15841  
 15901  
 15961  
 16021  
 16081  
 16141  
 16201  
 16261  
 16321  
 16381  
 16441  
 16501  
 16561  
 16621  
 16681  
 16741  
 16801  
 16861  
 16921  
 16981  
 17041  
 17101  
 17161  
 17221  
 17281  
 17341  
 17401  
 17461  
 17521  
 17581  
 17641  
 17701  
 17761  
 17821  
 17881  
 17941  
 18001  
 18061  
 18121  
 18181  
 18241  
 18301  
 18361  
 18421  
 18481  
 18541  
 18601  
 18661  
 18721  
 18781  
 18841  
 18901  
 18961  
 19021  
 19081  
 19141  
 19201  
 19261  
 19321  
 19381  
 19441  
 19501  
 19561  
 19621  
 19681  
 19741  
 19801  
 19861  
 19921  
 19981  
 20041  
 20101  
 20161  
 20221  
 20281  
 20341  
 20401  
 20461  
 20521  
 20581  
 20641  
 20701  
 20761  
 20821  
 20881  
 20941  
 21001  
 21061  
 21121  
 21181  
 21241  
 21301  
 21361  
 21421  
 21481  
 21541  
 21601  
 21661  
 21721  
 21781  
 21841  
 21901  
 21961  
 22021  
 22081  
 22141  
 22201  
 22261  
 22321  
 22381  
 22441  
 22501  
 22561  
 22621  
 22681  
 22741  
 22801  
 22861  
 22921  
 22981  
 23041  
 23101  
 23161  
 23221  
 23281  
 23341  
 23401  
 23461  
 23521  
 23581  
 23641  
 23701  
 23761  
 23821  
 23881  
 23941  
 24001  
 24061  
 24121  
 24181  
 24241  
 24301  
 24361  
 24421  
 24481  
 24541  
 24601  
 24661  
 24721  
 24781  
 24841  
 24901  
 24961  
 25021  
 25081  
 25141  
 25201  
 25261  
 25321  
 25381  
 25441  
 255

40 In Table 5 bold lines demarcate the individual oligonucleotides. Segment M contains 252 base pairs and has destroyed EcoRI, restriction sites at both ends. (Additional restriction sites within segment M are indicated). Segment M is inserted into vector pIC20R at an EcoRI restriction site and cloned.

As proposed in Figure 3, segment M is joined to segment A in the plasmid in which it is contained. Segment M is excised at the flanking restrictions sites from its cloning vector and spliced into pIC20K, harboring segment A, through successive digestions with HindIII followed by BglII. The pIC20K vector now comprises segment A joined to segment M with a HindIII site at the splice site (see Figure 3). Plasmid pIC20K is derived from pIC20R by removing the ScaI-NdeI DNA fragment and inserting a HincII fragment containing an NPTI coding region. The resulting plasmid of 4.44 kb confers resistance to kanamycin on E. coli.

The synthetic Btt gene is designed so that it is expressed in the pIC20R-kan vector in which it is constructed. This expression is produced utilizing the initiation methionine of the lacZ protein of pIC20K. The wild-type Btt crystal protein sequence expressed in this manner has full insecticidal activity. In addition, the synthetic gene is designed to contain a BamHI site 5' proximal to the initiating methionine codon and a BglII site 3' to the terminal TAG translation stop codon. This facilitates the cloning of the insecticidal crystal protein coding region into bacterial expression vectors such as pDR540 (Russell and Bennett, 1982). Plasmid pDR540 contains the TAC promoter which allows the production of proteins including Btt crystal

protein under controlled conditions in amounts up to 10% of the total bacterial protein. This promoter functions in many gram-negative bacteria including *E. coli* and *Pseudomonas*.

Production of Bt insecticidal crystal protein from the synthetic gene in bacteria demonstrates that the protein produced has the expected toxicity to coleopteran insects. These recombinant bacterial strains in themselves have potential value as microbial insecticides, product of the synthetic gene.

#### Example 4. Expression of a synthetic crystal protein gene in plants

The synthetic *Bt* crystal protein gene is designed to facilitate cloning into the expression cassettes. These utilize sites compatible with the *Bam*HI and *Bgl*II restriction sites flanking the synthetic gene. Cassettes are available that utilize plant promoters including CaMV 35S, CaMV 19S and the ORF 24 promoter from T-DNA. These cassettes provide the recognition signals essential for expression of proteins in plants. These cassettes are utilized in the micro Ti plasmids such as pH575. Plasmids such as pH575 containing the synthetic *Bt* gene directed by plant expression signals are utilized in disarmed *Agrobacterium tumefaciens* to introduce the synthetic gene into plant genomic DNA. This system has been described previously by Adang et al. (1987) to express *Bt* var. *kurstaki* crystal protein gene in tobacco plants. These tobacco plants were toxic to feeding tobacco hornworms.

#### Example 5. Assay for insecticidal activity

Bioassays were conducted essentially as described by Sekar, V. et al. *supra*. Toxicity was assessed by an estimate of the LD<sub>50</sub>. Plasmids were grown in *E. coli* JM105 (Yanisch-Perron, C. et al. (1985) *Gene* 33:103-119). On a molar basis, no significant differences in toxicity were observed between crystal proteins encoded by p544Pst-Met5, p544-HindIII, and pNSBP544. When expressed in plants under identical conditions, cells containing protein encoded by the synthetic gene were observed to be more toxic than those containing protein encoded by the native *Bt* gene. Immunoblots ("western" blots) of cell cultures indicated that those that were more toxic had more crystal protein antigen. Improved expression of the synthetic *Bt* gene relative to that of a natural *Bt* gene was seen as the ability to quantitate specific mRNA transcripts from expression of synthetic *Bt* genes on Northern blot assays.

#### Claims

1. A modified *Bt* insecticidal protein gene, the coding region of which has an A + T content no greater than 60%.
2. A modified *Bt* insecticidal protein gene according to claim 1 which encodes a protein which is at least 75% homologous to a native insecticidal protein of *Bt*.
3. A modified *Bt* insecticidal protein gene according to claim 1 which is at least about 85% homologous to a native insecticidal protein gene of *Bt*.
4. A modified *Bt* insecticidal protein gene according to claim 1 having the DNA sequence presented in Figure 1, spanning nucleotides 1 through 1793.
5. A modified *Bt* insecticidal protein gene according to claim 1 having the DNA sequence presented in Figure 1 spanning nucleotides 1 through 1833.
6. A modified *Bt* insecticidal protein gene according to claim 1 wherein the A + T content of said coding region is approximately 55%.
7. A modified *Bt* insecticidal protein gene according to claim 1 wherein a plant initiation sequence is present at the 5' end of the coding region.
8. A modified *Bt* insecticidal protein gene according to claim 1 wherein plant polyadenylation signals, comprising those having AATAAA, AATGAA, AATAAT, AATATT, GATAAA and AATAAG motifs, are eliminated in said DNA sequence.

9. A modified Bt insecticidal protein gene according to claim 1 wherein the polymerase II termination sequence, CAN<sub>7-9</sub>AGTNNA, is eliminated in said DNA sequence.
10. A modified Bt insecticidal protein gene according to claim 1 wherein CUUCGG hairpins are eliminated in said DNA sequence.
11. A modified Bt insecticidal protein gene according to claim 1 wherein plant consensus splice sites, including 5' = AAG:GTAAGT and 3' = TTTT(Pu)TTT(Pu)T(Pu)T(Pu)TGCAG:C, are eliminated in said DNA sequence.
12. A recombinant DNA cloning vector comprising a modified Bt insecticidal protein gene according to any preceding claim.
13. A plant cell which contains a modified Bt insecticidal protein gene according to any of claims 1 to 12.
14. A plant cell containing a DNA sequence encoding an insecticidal protein, said DNA sequence being derived from a Bt insecticidal protein gene, characterised in that the DNA sequence of the coding region of the Bt gene has been modified so as to have an A + T content no greater than 60%.
15. A plant cell according to claim 14, wherein said A + T content is approximately 55%.
16. A dicot plant cell according to any of claims 13 to 15, wherein the ratio of XCG codons to XCC codons in said modified gene (X being A, T, C or G) is approximately 0.3:1, and the ratio of XTA to XTT codons in said synthetic gene is approximately 0.35:1.
17. A monocot plant cell according to any of claims 13 to 15, wherein the ratio of XCG codons to XCC codons in said modified gene (X being A, T, C or G) is approximately 0.61:1, and the ratio of XTA to XTT codons in said synthetic gene is approximately 0.47:1.
18. A maize plant cell according to claim 13.
19. A method of producing a protein toxic to an insect, said method comprising maintaining a plant cell according to any of claims 13 to 18 or progeny thereof under conditions which cause said modified gene to be expressed in said plant cell or in said progeny thereof.
20. A method of producing a gene encoding an insecticidal protein, said method comprising:
  - (a) analyzing the coding sequence of a Bt insecticidal protein gene,
  - (b) modifying said coding sequence so as to have an A + T content no greater than 60%, and
  - (c) preparing DNA having said modified coding sequence.
21. A method according to claim 20 further comprising the step of modifying a portion of said coding sequence to eliminate CUUCGG hairpins.
22. A method according to claim 20 further comprising the step of modifying a portion of said coding sequence to eliminate plant polyadenylation signals including AATAAA, AATGAA, AATAAT, AATATT, GATAAA, and AATAAG.
23. A method according to claim 20 further comprising the step of modifying a portion of said coding sequence to eliminate polymerase II termination sequences, including CAN<sub>7-9</sub>AGTNNA.
24. A method according to claim 20 further comprising the step of modifying a portion of said coding sequence to eliminate plant consensus splice sites, including 5' = AAG:GTAAGT and 3' = TTTT(Pu)TTT(Pu)T(Pu)T(Pu)TGCAG:C.
25. A method according claim 20 further comprising the step of modifying a portion of said coding sequence to yield a sequence containing a plant translation initiation sequence.

26. A method according to claim 20 wherein said synthetic gene is intended for expression in dicot plants, said method further comprising the step of modifying a portion of said coding sequence to yield a ratio of XGC to XCC codons (X being A, T, C or G) of approximately 0.3:1 and a ratio of XTA to XTT codons of approximately 0.35:1.

5

27. A method according to claim 20 wherein said synthetic gene is intended for expression in monocot plants, said method further comprising the step of modifying a portion of said coding sequence to yield a ratio of XGC to XCC codons (X being A, T, C or G) of approximately 0.61:1 and a ratio of XTA to XTT codons of approximately 0.47:1.

10

28. A method according to any of claims 20 to 27, comprising the further steps of introducing said DNA into a host plant cell and reproducing said host plant cell containing said DNA.

### Patentansprüche

15

1. Modifiziertes Bt-Gen für ein insektizides Protein, dessen codierende Region einen A+T-Gehalt von höchstens 60% aufweist.

20

2. Modifiziertes Bt-Gen für ein insektizides Protein nach Anspruch 1, das für ein Protein codiert, das zu mindestens 75% mit einem nativen insektiziden Protein von Btt homolog ist.

3. Modifiziertes Bt-Gen für ein insektizides Protein nach Anspruch 1, das zu mindestens 85% mit einem nativen Gen von Btt für ein insektizides Protein homolog ist.

25

4. Modifiziertes Bt-Gen für ein insektizides Protein nach Anspruch 1, das die in Figur 1 dargestellte, sich über die Nucleotide 1 bis 1793 erstreckende DNA-Sequenz aufweist.

5. Modifiziertes Bt-Gen für ein insektizides Protein nach Anspruch 1, das die in Figur 1 dargestellte, sich über die Nucleotide 1 bis 1833 erstreckende DNA-Sequenz aufweist.

30

6. Modifiziertes Bt-Gen für ein insektizides Protein nach Anspruch 1, worin der A+T-Gehalt der codierenden Region etwa 55% beträgt.

35

7. Modifiziertes Bt-Gen für ein insektizides Protein nach Anspruch 1, worin sich an dem 5'-Ende der codierenden Region eine pflanzliche Initiationssequenz befindet.

8. Modifiziertes Bt-Gen für ein insektizides Protein nach Anspruch 1, worin pflanzliche Polyadenylierungssignale, jene mit AATAAA-, AATGAA-, AATAAT-, AATATT-, GATAAA- und AATAAG-Motiven eingeschlossen, in der DNA-Sequenz eliminiert sind.

40

9. Modifiziertes Bt-Gen für ein insektizides Protein nach Anspruch 1, worin die Polymerase-II-Terminationssequenz CAN<sub>7-9</sub>AGTNNA in der DNA-Sequenz eliminiert ist.

45

10. Modifiziertes Bt-Gen für ein insektizides Protein nach Anspruch 1, worin CUUCGG-Haarnadelschleifen in der DNA-Sequenz eliminiert sind.

11. Modifiziertes Bt-Gen für ein insektizides Protein nach Anspruch 1, worin pflanzliche Consensus-Spleißstellen, einschließlich 5' = AAG:GTAAGT und 3' = TTTT(Pu)TTT(Pu)T(Pu)T(Pu)TGCAG:C, in der DNA-Sequenz eliminiert sind.

50

12. Rekombinanter DNA-Klonierungsvektor, der ein modifiziertes Bt-Gen für ein insektizides Protein nach einem der vorhergehenden Ansprüche umfaßt.

55

13. Pflanzenzelle, die ein modifiziertes Bt-Gen für ein insektizides Protein nach einem der Ansprüche 1 bis 12 enthält.

14. Pflanzenzelle, die eine DNA-Sequenz enthält, welche für ein insektizides Protein codiert, wobei diese DNA-Sequenz aus einem Bt-Gen für ein insektizides Protein stammt, dadurch gekennzeichnet, daß die

DNA-Sequenz der codierenden Region des Bt-Gens so modifiziert worden ist, daß sie einen A + T-Gehalt von höchstens 60% aufweist.

15. Pflanzenzelle nach Anspruch 14, worin der A + T-Gehalt etwa 55% beträgt.
- 5 16. Zelle einer zweikeimblättrigen Pflanze nach einem der Ansprüche 13 bis 15, worin das Verhältnis von XCG-Codons zu XCC-Codons in dem modifizierten Gen (wobei X A, T, C oder G ist) etwa 0,3:1 ist und das Verhältnis von XTA- zu XTT-Codons in dem synthetischen Gen etwa 0,35:1 ist.
- 10 17. Zelle einer einkeimblättrigen Pflanze nach einem der Ansprüche 13 bis 15, worin das Verhältnis von XCG-Codons zu XCC-Codons in dem modifizierten Gen (wobei X A, T, C oder G ist) etwa 0,61:1 ist und das Verhältnis von XTA- zu XTT-Codons in dem synthetischen Gen etwa 0,47:1 ist.
18. Maispflanzenzelle nach Anspruch 13.
- 15 19. Verfahren zur Erzeugung eines Proteins, das für ein Insekt toxisch ist, wobei dieses Verfahren umfaßt, daß eine Pflanzenzelle nach einem der Ansprüche 13 bis 18 oder deren Nachkommen unter Bedingungen gehalten werden, welche die Expression des modifizierten Gens in dieser Pflanzenzelle oder in deren Nachkommen bewirken.
- 20 20. Verfahren zur Herstellung eines Gens, das für ein insektizides Protein codiert, wobei dieses Verfahren umfaßt:
  - (a) das Analysieren der codierenden Sequenz eines Bt-Gens für ein insektizides Protein,
  - (b) das Modifizieren der codierenden Sequenz, so daß sie einen A + T-Gehalt von höchstens 60%
  - 25 aufweist, und
  - (c) die Herstellung von DNA mit dieser modifizierten codierenden Sequenz.
21. Verfahren nach Anspruch 20, das weiters den Schritt der Modifizierung eines Teils der codierenden Sequenz umfaßt, um CUUCGG-Haarnadelschleifen zu eliminieren.
- 30 22. Verfahren nach Anspruch 20, das weiters den Schritt der Modifizierung eines Teils der codierenden Sequenz umfaßt, um pflanzliche Polyadenylierungssignale, einschließlich AATAAA, AATGAA, AATAAT, AATATT, GATAAA und AATAAG, zu eliminieren.
- 35 23. Verfahren nach Anspruch 20, das weiters den Schritt der Modifizierung eines Teils der codierenden Sequenz umfaßt, um Polymerase-II-Terminationssequenzen, einschließlich CAN<sub>7-9</sub>AGTNNAA, zu eliminieren.
- 40 24. Verfahren nach Anspruch 20, das weiters den Schritt der Modifizierung eines Teils der codierenden Sequenz umfaßt, um pflanzliche Consensus-Spleißstellen, einschließlich 5' = AAG:GTAAGT und 3' = TTTT(Pu)TTT(Pu)T(Pu)T(Pu)TGCAG:C, zu eliminieren.
- 45 25. Verfahren nach Anspruch 20, das weiters den Schritt der Modifizierung eines Teils der codierenden Sequenz umfaßt, um eine Sequenz zu erhalten, die eine pflanzliche Translations-Initiationssequenz enthält.
26. Verfahren nach Anspruch 20, worin das synthetische Gen für die Expression in zweikeimblättrigen Pflanzen bestimmt ist, wobei dieses Verfahren weiters den Schritt der Modifizierung eines Teils der codierenden Sequenz umfaßt, um ein Verhältnis von XGC- zu XCC-Codons (wobei X A, T, C oder G
- 50 ist) von etwa 0,3:1 und ein Verhältnis von XTA- zu XTT-Codons von etwa 0,35:1 zu erhalten.
27. Verfahren nach Anspruch 20, worin das synthetische Gen für die Expression in einkeimblättrigen Pflanzen bestimmt ist, wobei dieses Verfahren weiters den Schritt der Modifizierung eines Teils der codierenden Sequenz umfaßt, um ein Verhältnis von XGC- zu XCC-Codons (wobei X A, T, C oder G
- 55 ist) von etwa 0,61:1 und ein Verhältnis von XTA- zu XTT-Codons von etwa 0,47:1 zu erhalten.
28. Verfahren nach einem der Ansprüche 20 bis 27, das die weiteren Schritte der Einführung der DNA in eine Wirtspflanzenzelle und der Reproduktion der diese DNA enthaltenden Wirtspflanzenzelle umfaßt.

## Revendications

1. Gène modifié d'une protéine insecticide Bt, dont la région codante possède une teneur en A + T ne dépassant pas 60%.
2. Gène modifié d'une protéine insecticide Bt selon la revendication 1, qui code pour une protéine qui est au moins à 75% homologue à une protéine insecticide d'origine naturelle de Btt.
3. Gène modifié d'une protéine insecticide Bt selon la revendication 1, qui est au moins à environ 85% homologue à un gène de protéine insecticide d'origine naturelle de Btt.
4. Gène modifié d'une protéine insecticide Bt selon la revendication 1 ayant la séquence ADN présentée dans la figure 1, portant sur les nucléotides de 1 à 1793.
5. Gène modifié d'une protéine insecticide Bt selon la revendication 1 ayant la séquence ADN présentée dans la figure 1 portant sur les nucléotides de 1 à 1893.
6. Gène modifié d'une protéine insecticide Bt selon la revendication 1, dans lequel la teneur en A + T de ladite région codante est approximativement de 55%.
7. Gène modifié d'une protéine insecticide Bt selon la revendication 1, dans lequel une séquence d'initiation végétale est présente à la terminaison 5' de la région codante.
8. Gène modifié d'une protéine insecticide Bt selon la revendication 1, dans lequel les signaux de polyadénylation végétaux, comprenant ceux ayant les motifs AATAAA, AATGAA, AATAAT, AATATT, GATAAA et AATAAG, sont éliminés dans ladite séquence ADN.
9. Gène modifié d'une protéine insecticide Bt selon la revendication 1, dans lequel la séquence de terminaison de polymérase II, CAN<sub>7-9</sub>AGTN<sub>10-12</sub>AA, est éliminée dans ladite séquence ADN.
10. Gène modifié d'une protéine insecticide Bt selon la revendication 1, dans lequel les épingles à cheveux CUUCGG sont éliminées dans ladite séquence ADN.
11. Gène modifié d'une protéine insecticide Bt selon la revendication 1, dans lequel les sites d'épissage consensus végétaux, incluant 5' = AAG:GTAAGT et 3' = TTTT(Pu)TTT(Pu)T(Pu)T(Pu)TGCAG:C, sont éliminés dans ladite séquence ADN.
12. Vecteur clonant d'ADN recombinant comprenant un gène modifié d'une protéine insecticide Bt selon l'une quelconque des revendications précédentes.
13. Cellule végétale qui contient un gène modifié d'une protéine insecticide Bt selon l'une quelconque des revendications 1 à 12.
14. Cellule végétale contenant une séquence ADN codant pour une protéine insecticide, ladite séquence ADN étant dérivée d'un gène d'une protéine insecticide Bt, caractérisée en ce que la séquence ADN de la région codante du gène Bt a été modifiée de façon à avoir une teneur en A + T ne dépassant pas 60%.
15. Cellule végétale selon la revendication 14, dans laquelle ladite teneur en A + T est approximativement de 55%.
16. Cellule de plante dicotylédone selon l'une quelconque des revendications 13 à 15, dans laquelle le rapport de codons XCG aux codons XCC dans ledit gène modifié (X étant A, T, C ou G) est approximativement de 0,3:1, et le rapport des codons XTA aux codons XTT dans ledit gène synthétique est approximativement de 0,35:1.
17. Cellule de plante monocotylédone selon l'une quelconque des revendications 13 à 15, dans laquelle le rapport des codons XCG aux codons XCC dans ledit gène modifié (X étant A, T, C, ou G) est



approximativement de 0,61:1, et le rapport des codons XTA aux codons XTT dans ledit gène synthétique est approximativement de 0,47:1.

18. Cellule végétale de maïs selon la revendication 13.

19. Procédé pour produire une protéine toxique envers un insecte, ledit procédé consistant à maintenir une cellule végétale selon l'une quelconque des revendications 13 à 18 ou sa descendance dans des conditions qui entraînent ledit gène modifié à s'exprimer dans ladite cellule végétale ou dans la descendance de celle-ci.

20. Procédé pour produire un gène codant pour une protéine insecticide, ledit procédé consistant:  
 (a) à analyser la séquence codante d'un gène d'une protéine insecticide Bt,  
 (b) à modifier ladite séquence codante de façon à avoir une teneur en A + T ne dépassant pas 60%,  
 et  
 (c) à préparer l'ADN ayant ladite séquence codante modifiée.

21. Procédé selon la revendication 20, comprenant de plus l'étape consistant à modifier une portion de ladite séquence codante pour éliminer les épingles à cheveux CUUCGG.

22. Procédé selon la revendication 20 comprenant de plus l'étape consistant à modifier une portion de ladite séquence codante pour éliminer les signaux végétaux de polyadénylation incluant AATAAA, AATGAA, AATAAT, AATATT, GATAAA et AATAAG.

23. Procédé selon la revendication 20, comprenant de plus l'étape consistant à modifier une portion de ladite séquence codante pour éliminer les séquences de terminaison de la polymérase II, incluant CAN<sub>7-9</sub>AGTNNA.

24. Procédé selon la revendication 20, comprenant de plus l'étape consistant à modifier une portion de ladite séquence codante pour éliminer les sites végétaux d'épissage consensus, incluant 5' = AAG:GTAAGT et 3' = TTTT(Pu)TTT(Pu)T(Pu)T(Pu)TGACG:C.

25. Procédé selon la revendication 20 comprenant de plus l'étape consistant à modifier une portion de ladite séquence codante pour obtenir une séquence contenant une séquence végétale d'initiation de la traduction.

26. Procédé selon la revendication 20, dans lequel ledit gène synthétique est prévu pour s'exprimer dans des plantes dicotylédones, ledit procédé comprenant de plus l'étape consistant à modifier une portion de ladite séquence codante pour produire un rapport des codons XGC aux codons XCC (X étant A, T, C ou G) d'approximativement 0,3:1 et un rapport des codons XTA aux codons XTT d'approximativement 0,35:1.

27. Procédé selon la revendication 20, dans lequel le gène synthétique est prévu pour s'exprimer dans des plantes monocotylédones, ledit procédé comprenant de plus l'étape consistant à modifier une portion de ladite séquence codante pour obtenir un rapport des codons XGC aux codons XCC (X étant A, T, C ou G) d'approximativement 0,61:1 et un rapport des codons XTA aux codons XTT d'approximativement 0,47:1.

28. Procédé selon l'une quelconque des revendications 20 à 27, comprenant les étapes supplémentaires consistant à introduire ledit ADN dans une cellule végétale hôte et à reproduire ladite cellule végétale hôte contenant ledit ADN.

FIGURE 1

A T T T A A A A A A T A A A  
 1 ATGGCTGCAGACAACCAACGGAGGCCCTCGATAGCTCTACCAACCAAGATGTCATTGAGAGGGCATCTCCGTGTGGGTGATCTCCTTGGCGTGTG 100  
 M A A D M N T E A L D S S T T K O V I O K G I S V V G D L L G V V G  
 G A G T T A T A A T G  
 101 GTTCCCTCTTGGTGGTGGCCCTTGTTCGTCTACACTAATCTTCTGAATACTATTGGCCAGCGAAGACCCCTTGAAGGCTTTTATGGAGCAAGTGA 200  
 F P F G G A L V S F Y T N F L N T I U P S E D P V K A F M E Q V E  
 A A A A T A A T T  
 201 AGCTTTGATGGATCAGAAGATCGCTGATTATGCAAGAACAAGCTCTTCTGAGCTCCAGGGCTTCAGAACACGTCGAAGATTATGTGAGTGCACTG 300  
 A L M D O K I A D Y A K N K A L A E L Q G L O N H V E D Y V S A L  
 A A G T G A T T T T  
 301 AGTTCATGGCAAAAGATCCTGTGTCTCAGCAATCCACATAGCCAGGGGGCATAAGGGAGCTGTTCTCTCAAGCAGAAAGTCACCTTCGGGAATTCAA 400  
 S S W Q K M P V S S R N P H S Q G R I R E L F S Q A E S H F R N S H  
 G A T A A A A T A T  
 401 TGCCCTCCTTGGCATCTCTGGGTACGAGGTCTCTTCTTCTACAACTACGCTCAAGCTGCCAACACACATCTGTTCTTAAAGAGGCTCAAACTCA 500  
 P S F A I S G Y E V L F L T T Y A Q A A N T H L F L L K D A Q I Y  
 A A A T A A A T  
 501 TGGTGAAGAATGGGATACGAGAAGAAGATACGCTGAGTCTACAGGGCTCAACTAAACTTACTCAAGCATATACTGACCACTGTGTCAATGGTAT 600  
 G E E V G Y E K E D I A E F Y K R O L K L T Q E Y T O M C V K W Y  
 A A A T T A A T A T A  
 601 AATGTTGATTCGATAAGTTGAGAGGTTTCATCTTATGAATCTTGGGTAACTTAAACCGGTACCGCAGAGAGATGACATTGACAGTCTGCACTTGTAT 700  
 N V G L D K L R G S S Y E S V M V M F N R Y R R E N T L T V L D L I A  
 G A A A A A  
 701 CACTATTTCATGTATGATGTTCCACTACCCAAAGAGGTTAAACCGAATGACTAGAGACCTTTAAACCGATCCCATTTGCGGAGTCAACACCT 800  
 L F P L Y D V R L Y P K E V K T E L T R D V L T D P I V G V N H L  
 T G T T T A T A T A  
 801 TGGGTACGGCAACACCTTCTCTAATAGAAAACATACCTGTAACACCATCTATTCGACTATCTGCACAGAATTCAGTTTCACACGGGTTCAC 900  
 R G Y G T T F S M I E N Y I R K P N L F D Y L H R I O F H T R F Q  
 T T A A T A A  
 901 CCAGGATCTATGCAATGACTCTTTCAACTATTGGTCCGGTAATTATGTTTCACTAGACCCAGCATAGGATCTAATGACATCATCAGCTCTCCATTCT 1000  
 P G Y Y G N D S F N Y M S G N Y V S T R P S I G S M D I I T S P F Y  
 T T A A G T A A T A A A A A G  
 1001 ACCGAAACAGTCTCCGAGCTGTGCAAACTTGGAGTTAATGGAGAGAAGTCTATAGAGCGGTGCCAATACCAATCTTGGCGTCTGGCGTCCGC 1100  
 G N K S S E P V O N L E F N G E K V Y T R A V A N T N L A V U P S A  
 A T A A A A A A  
 1101 TGGTACTCAGGTGTTACCAAGTGAATTCAGCAATACATGATCAGACAGATCAAGCAAGTACTCAACTTACCACTCAAGAGGAATGTTGGCGCG 1200  
 V Y S G V T K V E F S O Y N D Q T D E A S T Q T Y D S K R N V G A  
 T G A A A A T  
 1201 GTCAGCTGGATCTATCGATCACTCCCTCCAGAACCCAGGATGAACCTCTAGACAAGGCTTATAGCCATCACTCAATTACGTAATGTCTTCTCA 1300  
 V S V D S I D G L P P E T T D E P L E K G Y S H O L N Y V N C F L M  
 A A A T A T A A  
 1301 TGCAGGGTAGTAGAGGTACCATCCAGTGTAACTTGGACTCACAGAGTGTAGACTTCTTCAACATGATTGATTCGAAAAAGATTACTCAACTCCGT 1400  
 Q G S R G T I P V L T M T H K S V D F F N M I D S K K I T O L P L  
 A A A A A A A A G T  
 1401 CGTAAGGCCCTACAAGTTACAATCGTCTCCCTTGTCCGAGTCTAGCTTACAGGAGCAGATATCATTCAATGCACTGAGAATGGGTCCGCGGCA 1500  
 V K A Y K L Q S G A S V V A G P R F T G G D I I O C T E N G S A A  
 T G A A A A T C T A  
 1501 ACTATCTACGTACACCTGATGTGTCTACTCTCAAAAGTATCGTGTAGAAATCATTATGCTTCTACCTCTCAGATAACATTCACTAAGCTTGGAGG 1600  
 T I Y V T P D V S Y S Q K Y R A R I N Y A S T S O I T F T L S L D G  
 A T T I A G A T T A T A T A T  
 1601 GGGCTCCATTCAACCAATACTACTTGGATAAGACCATCAACAAGGAGACACACTCAGGTATAATTCATTCAACTAGCCAGCTTCAGCACTCCATTCCA 1700  
 A P F N Q Y T F D K T I N K G D T L T Y N S F N L A S F S T P F E  
 A T A A A A A A A T T A A  
 1701 ATTGTCAAGGACCACTTCCAGATAGGCTCAGAGGATGAGTGTGGTGGTACAGGTTACATCGACAAGATTGAGTTCATTCCAGTCAACCTTAGGCTC 1800  
 L S G N H L O I G V T G L S A G D K V Y I D K I E F I P V M L R S  
 CCAGGAACCGAGCTTGAGTTTCATCGACATCTAG  
 1801 P G T E L E F I D I \*


Figure 2. Simplified Scheme for Synthesis of Btt Gene

FIGURE 3

